

10-18-00
534 Rec'd PCT/PTO 16 OCT 2000

FORM PTO-1390		U S DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE	ATTORNEY'S DOCKET NUMBER 245-55928
TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. § 371			U S APPLICATION NO (If known, see 37 C F R § 1.5) 09/673763
INTERNATIONAL APPLICATION NO. PCT/US99/08744	INTERNATIONAL FILING DATE 20 April 1999	PRIORITY DATE CLAIMED 20 April 1998	
TITLE OF INVENTION CHLAMYDIA PROTEINS AND THEIR USES			
APPLICANT(S) FOR DO/EO/US Rockey and Bannantine			
Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information			
<ol style="list-style-type: none"> <input checked="" type="checkbox"/> This is a FIRST submission of items concerning a filing under 35 U.S.C. § 371. <input type="checkbox"/> This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. § 371 <input type="checkbox"/> This express request to begin national examination procedures (35 U.S.C. § 371(f) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. § 371(b) and PCT Articles 22 and 39(1) <input checked="" type="checkbox"/> A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date <input checked="" type="checkbox"/> A copy of the International Application as filed (35 U.S.C. § 371(c)(2)) <ol style="list-style-type: none"> <input checked="" type="checkbox"/> is transmitted herewith (required only if not transmitted by the International Bureau). <input type="checkbox"/> has been transmitted by the International Bureau. <input type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US) <input type="checkbox"/> A translation of the International Application into English (35 U.S.C. § 371(c)(2)) <input checked="" type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. § 371(c)(3)) <ol style="list-style-type: none"> <input type="checkbox"/> are transmitted herewith (required only if not transmitted by the International Bureau). <input type="checkbox"/> have been transmitted by the International Bureau <input type="checkbox"/> have not been made; however, the time limit for making such amendments has NOT expired. <input checked="" type="checkbox"/> have not been made and will not be made <input type="checkbox"/> A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. § 371(c)(3)). <input checked="" type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. § 371(c)(4)). <input type="checkbox"/> A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. § 371(c)(5)) 			
Items 11. to 16. below concern document(s) or information included:			
<ol style="list-style-type: none"> <input checked="" type="checkbox"/> An Information Disclosure Statement under 37 C.F.R. §§ 1.97 and 1.98, including copies of references cited in International Search Report. <input checked="" type="checkbox"/> An assignment document for recording. A separate cover sheet in compliance with 37 C F R. §§ 3.28 and 3.31 is included <input checked="" type="checkbox"/> A FIRST preliminary amendment. <ol style="list-style-type: none"> <input type="checkbox"/> A SECOND or SUBSEQUENT preliminary amendment <input type="checkbox"/> A substitute specification <input type="checkbox"/> A change of power of attorney and/or address letter. <input checked="" type="checkbox"/> Other items or information. <ol style="list-style-type: none"> <input checked="" type="checkbox"/> Written Opinion. <input type="checkbox"/> Preliminary Examination Report. <input checked="" type="checkbox"/> International Search Report <input checked="" type="checkbox"/> Computer Readable Form of Sequence Listing. <input checked="" type="checkbox"/> Statement in Compliance with 37 C F R § 1.821(f) 			



24197

PATENT

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of: Rockey and Bannantine

Art Unit: Not yet assigned

Application No.: Not yet assigned

CERTIFICATE OF MAILING


Filed: Herewith

I hereby certify that this paper and the documents referred to as being attached or enclosed herewith are being deposited with the United States Postal Service on October 16, 2000 via Express Mail in an envelope addressed to: BOX PATENT APPLICATION, COMMISSIONER FOR PATENTS, WASHINGTON, D.C. 20231.

For: CHLAMYDIA PROTEINS AND THEIR USES

Examiner: Not yet assigned

Date: October 16, 2000


 Tanya M. Harding, Ph.D.
 Attorney for Applicant

BOX PATENT APPLICATION
 COMMISSIONER FOR PATENTS
 WASHINGTON, D.C. 20231

PRELIMINARY AMENDMENT

Please enter the following amendment in the above-referenced patent application:

In the Specification:

At page 9, line 35, please replace the term "DH5□" with --DH5α--.

At page 12, line 12, please replace the term "DH5□" with --DH5α--.

At page 13, line 35, please replace the term "DH5□" with --DH5α--.

At page 25, please insert the following complete citation after line 8: --Bannantine, J.P., *et al.* (1998) *Infect. Immun.* 66:6017-6021.--.

REMARKS

By this voluntary amendment, the Specification of the application has been amended solely to correct administrative and typographical errors and omissions. The *E. coli* strain DH5α is well known, and the corrections to this term are made only to correct a printing error. Support for the insertion of the Bannantine *et al.* (1998) citation can be found at page


PATENT

14, line 17, which provides sufficient information to locate this article. Applicants believe this amendment adds no new matter to the application.

Respectfully submitted,

KLARQUIST SPARKMAN CAMPBELL
LEIGH & WHINSTON, LLP

By



Tanya M. Harding, Ph.D.
Registration No. 42,630

One World Trade Center, Suite 1600
121 S.W. Salmon Street
Portland, Oregon 97204
Telephone: (503) 226-7391
Facsimile: (503) 228-9446

09/673763

CHLAMYDIA PROTEINS AND THEIR USES

I. FIELD OF THE INVENTION

The present invention relates to the detection of *Chlamydia* and to the diagnosis, treatment
5 and prevention of *Chlamydia* infections in animals.

II. BACKGROUND

Chlamydiae are obligate intracellular bacterial pathogens with a unique biphasic life cycle. They appear as two distinct cellular types, a small dense cell or elementary body (EB) that is
10 enclosed in a rigid bacterial cell wall, and a larger metabolically active reticulate body (RB). The EB is resistant to physical disruption and is infectious, whereas the RB is more fragile and only exists inside cells. The *Chlamydia* life cycle begins with the attachment of the EB form to the host cell which is followed by endocytosis into a nascent vacuole, also called an "inclusion membrane."

After EB attachment and entry, replication of the EB form produces RB forms that continue to
15 grow within the vacuole. By 72 hour post-infection, this growth phase is terminated when the RBs condense, and reorganize back to EBs. The lysis of the host cell results in release of EBs to infect new host cells. The difficulties in working with *Chlamydiae* center on the obligate intracellular requirement for growth and the fact that no adequate genetic engineering methods have been developed for this organism.

The genus *Chlamydia* includes two species that are primarily associated with human
20 disease: *C. trachomatis* and *C. pneumoniae*. *C. trachomatis* causes trachoma, an eye disease that is the leading cause of preventable infectious blindness worldwide with an estimated 500 million cases of active trachoma worldwide. *C. trachomatis* also causes a sexually transmitted chlamydial disease which is very common worldwide. *C. trachomatis* also causes lymphogranuloma
25 venereum, a debilitating systemic disease characterized by lymphatic gland swelling. The most serious sequelae of chlamydial genital infections of females include salpingitis, pelvic inflammatory disease, and ectopic pregnancy. In the US alone, it is estimated that over 4 million new sexually transmitted *C. trachomatis* infections occurred in 1990, leading to over four billion dollars in direct and indirect medical expenses. The World Health Organization estimates that 89
30 million new cases of genital *Chlamydia* occurred worldwide in 1995 (Peeling and Brunham, 1996).

C. pneumoniae causes respiratory diseases including so called walking pneumonia, a low-
grade disease such that the infected person frequently fails to obtain treatment and remains in the community as an active, infectious carrier. *C. pneumoniae* is currently of interest because of its strong epidemiological association with coronary artery disease, and there is also some evidence to
35 link it with multiple sclerosis.

Of the other disease-causing species of *Chlamydia*, *Chlamydia psittaci* and *Chlamydia pecorum* are primarily pathogens of wild and domestic animals, but these species may infect

humans accidentally. *C. psittaci* is acquired through respiratory droplet infection and is considered an occupational health hazard for bird fanciers and poultry workers.

There is tremendous interest in the identification of candidate antigens for protection against chlamydial disease. While a prior infection with *C. trachomatis* will protect against a subsequent challenge by the same strain, indicating a protective component that stimulates the host immune response, most serious chlamydial diseases are exacerbated by an overaggressive anti-chlamydial immune response. Antigens recognized in the context of an infection appear to elicit a protective response whereas immunization with purified, killed (EB form) *Chlamydia* results in an immunopathological response. Therefore for the purposes of vaccine development, one needs to find epitopes that confer protection, but do not contribute to pathology. It is an object of this invention to provide *Chlamydia* polypeptides for use as vaccines that induce a protective immune response without inducing the pathological response caused by the antigens associated with the EB form of *Chlamydia*. Such immunostimulatory peptides will be useful in the treatment, as well as in the diagnosis, detection and prevention of Chlamydial infections.

III. SUMMARY OF THE INVENTION

The present invention includes the use of *Chlamydia* proteins that show enhanced expression in the reticulate body (RB) stage relative to the elementary body (EB) stage of the *Chlamydia* life cycle. These proteins are not present at detectable levels in the EB form using current immunological techniques and are thus said to be "infection-specific." Certain of these infection-specific proteins are found in the inclusion membrane of the infected cell, and so have been termed "Inc" proteins. These include the IncA, IncB, and IncC proteins of *Chlamydia* as described in the present disclosure. The genes that encode the IncA, IncB and IncC proteins are referred to as *incA*, *incB* and *incC* respectively. Other proteins of *Chlamydia* described herein have also been shown by the inventors to be infection-specific, but are not known to be incorporated into the inclusion membrane; these include the p242, TroA, and TroB proteins. The TroA and TroB proteins have been so named because they resemble the Tro proteins of *Treponema pallidum*, which are thought to form part of an ABC transport system.

The inventors have shown that the infection-specific *Chlamydia* proteins of the disclosure are recognized by convalescent antisera (i.e., antisera taken from an animal that has recovered from a *Chlamydia* infection) but are not recognized by antisera against the killed EB form of *Chlamydia*. Thus, the proteins are expressed only during active chlamydial infection and are therefore useful as protective antigens. These infection-specific proteins may be used to confer a protective immune response without inducing a pathological effect. Additionally, immunofluorescence microscopy and immunoblotting with antisera demonstrated that the infection-specific proteins are present in *Chlamydia*-infected HeLa cells, but are undetectable in purified EBs and absent in uninfected HeLa cells.

Immunofluorescence microscopy reveals that IncA, IncB and IncC are localized to the inclusion membrane of infected HeLa cells. Reverse-transcription polymerase chain reactions (RT-PCR), northern hybridization data, and restriction analysis revealed that the *incB* and *incC* genes are closely linked and transcribed in an operon. RT-PCR, restriction analysis and sequential Southern hybridizations of *incA* then *incC* to the same filter provided evidence that *incA* is separated from the *incB* and *incC* operon by about 110 kb. The *C. trachomatis* *Tro* genes are not closely linked with the p242 gene.

The present invention includes the nucleotide and amino acid sequences for certain infection-specific proteins from *Chlamydia*. These proteins are p242, TroA, and TroB from *C. trachomatis*, and the IncB, and IncC proteins from *C. psittaci*. The scope of the invention includes fragments of these proteins that may be used in a vaccine preparation or that may be used in a method of detecting *Chlamydia* antibodies. Such fragments may be, for example, 5, 10, 15, 20, 25, or 30 contiguous amino acids in length. They may even encompass the entire protein.

More specifically, the present invention encompasses the purified infection-specific proteins having amino acid sequences as shown in SEQ ID NOS: 2, 4, 6, 10, and 12, amino acid sequences that differ from such sequences by one or more conservative amino acid substitutions, and amino acid sequences that show at least 75% sequence identity with such amino acid sequences.

Then invention also includes isolated nucleic acid molecules that encode a protein as described in the above paragraph, including isolated nucleic acid molecules with nucleotide sequences as shown in SEQ ID NOS: 1,3, 5, 9, and 11.

The present invention also includes a vaccine or immunostimulatory preparation directed against the reticulate body (RB) form of *Chlamydia* comprising one or more purified infection-specific peptides (or portions or fragments thereof, or peptides showing sequence similarity to a portion of such a peptide). Such peptide fragments may be, for example, 5, 10, 15, 20, 25, or 30 contiguous amino acids in length, of the sequence shown in SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, or 18. Peptides used in such a vaccine may even encompass the entire purified peptide of SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, or 18, a peptide that differs from such a peptide by one or more conservative amino acid substitutions, or a peptide having at least 75% sequence identity with such a peptide. Such vaccine preparations may contain one or more pharmaceutically acceptable excipients, adjuvants or diluents.

The invention additionally encompasses methods for making a vaccine, comprising combining a pharmaceutically acceptable excipient with a peptide described herein. Also included is a method of vaccination comprising administering a vaccine as described herein to a mammal.

The present invention also provides a method for the diagnostic use of the disclosed purified infection-specific peptides, for instance by use in a diagnostic assay to detect the presence of infection-specific antibodies in a medical specimen, in which antibodies bind to the *Chlamydia* peptide and indicate that the subject from which the specimen was removed was previously

exposed to *Chlamydia*. Such a method may comprise: (i) supplying a biological sample, such as blood from an animal, that is suspected to contain infection-specific anti-*Chlamydia* antibody, (ii) contacting the sample with at least one infection-specific *Chlamydia* peptide described herein, such that a reaction between the peptide and the infection-specific anti-*Chlamydia* antibody gives rise to a detectable effect, such as a chromogenic conversion; and (iii) detecting this detectable effect.

The present invention also provides a method of using antibodies that bind specifically with the disclosed proteins for detection of infection-specific *Chlamydia* antigen, indicating the presence of *Chlamydia* in the RB stage as distinct from the EB stage. For instance, the relevant infection-specific antibodies may be used to provide specific binding in an Enzyme Linked Immunosorbant Assay (ELISA) or other immunological assay wherein the antibody F_c portion is linked to a chromogenic, fluorescent or radioactive molecule and the F_{ab} portion specifically interacts with, and binds to, an infection-specific protein. Such a method may comprise: (i) supplying a biological sample from an animal suspected to contain an infection-specific *Chlamydia* antigen, and (ii) contacting the sample with at least one infection-specific anti-*Chlamydia* antibody, such that a reaction between the antibody and the infection-specific *Chlamydia* protein gives rise to a detectable effect; and (iii) detecting this detectable effect.

Other aspects of the present invention include the use of probes and primers derived from the nucleotide sequences that encode infection-specific peptides, to detect the presence of *Chlamydia* nucleic acids in medical specimens. Such probes and primers may be nucleotide fragments, of, for example, 15, 20, 25, 30 or 40 contiguous nucleotides of the sequence shown in SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, or 17.

An additional aspect of the invention is a method of treating a *Chlamydia* infection by directing a therapeutic agent against a specific target, where the target is chosen from an infection specific protein of *Chlamydia*, a gene that encodes an infection-specific protein of *Chlamydia*, and an RNA transcript that encodes an infection-specific protein of *Chlamydia*, wherein the therapeutic agent interacts with said target to affect a reduction in pathology.

These and other aspects of the invention will become more apparent from the following description.

IV. SEQUENCE LISTING

SEQ ID NO:1 shows a nucleic acid sequence encoding the p242 *C. trachomatis* protein, with deduced primary amino acid sequence also shown.

SEQ ID NO:2 shows the amino acid sequence of the p242 *C. trachomatis* protein.

SEQ ID NO:3 shows a nucleic acid sequence encoding the TroA *C. trachomatis* protein, with deduced primary amino acid sequence also shown.

SEQ ID NO:4 shows the amino acid sequence of the TroA *C. trachomatis* protein.

SEQ ID NO:5 shows a nucleic acid sequence encoding the TroB *C. trachomatis* protein, with deduced primary amino acid sequence also shown.

SEQ ID NO:6 shows the amino acid sequence of the TroB *C. trachomatis* protein.

SEQ ID NO:7 shows a nucleic acid sequence encoding the IncA *C. psittaci* protein, with deduced primary amino acid sequence also shown.

SEQ ID NO:8 shows the amino acid sequence of the IncA *C. psittaci* protein.

5 SEQ ID NO:9 shows a nucleic acid sequence encoding the IncB *C. psittaci* protein, with deduced primary amino acid sequence also shown.

SEQ ID NO:10 shows the amino acid sequence of the IncB *C. psittaci* protein.

SEQ ID NO:11 shows a nucleic acid sequence encoding the IncC *C. psittaci* protein, with deduced primary amino acid sequence also shown.

10 SEQ ID NO:12 shows the amino acid sequence of the IncC *C. psittaci* protein.

SEQ ID NO:13 shows a nucleic acid sequence encoding the IncA *C. trachomatis* protein, with deduced primary amino acid sequence also shown.

SEQ ID NO:14 shows the amino acid sequence of the IncA *C. trachomatis* protein.

15 SEQ ID NO:15 shows a nucleic acid sequence encoding the IncB *C. trachomatis* protein, with deduced primary amino acid sequence also shown.

SEQ ID NO:16 shows the amino acid sequence of the IncB *C. trachomatis* protein.

SEQ ID NO:17 shows a nucleic acid sequence encoding the IncC *C. trachomatis* protein, with deduced primary amino acid sequence also shown.

SEQ ID NO:18 shows the amino acid sequence of the IncC *C. trachomatis* protein.

20 SEQ ID NO:19 shows the upstream oligonucleotide used to amplify the *C. psittaci* *incC* ORF.

SEQ ID NO:20 shows the downstream oligonucleotide used to amplify the *C. psittaci* *incC* ORF.

25 SEQ ID NO:21 shows the upstream oligonucleotide used to amplify the *C. psittaci* *incB* ORF.

SEQ ID NO:22 shows the downstream oligonucleotide used to amplify the *C. psittaci* *incB* ORF.

SEQ ID NO:23 shows the upstream oligonucleotide used to amplify the *C. psittaci* *incA* ORF.

30 SEQ ID NO:24 shows the downstream oligonucleotide used to amplify the *C. psittaci* *incA* ORF.

V. DESCRIPTION OF THE INVENTION

A. DEFINITIONS

35 Particular terms and phrases used herein have the meanings set forth below.

"EB" refers to the Elementary Body, an environmentally refractile and largely metabolically dormant form of *Chlamydia* that is infectious and is presented as a small dense body enclosed by a bacterial cell wall.

5 "RB" refers to the Reticulate Body, a metabolically active form of *Chlamydia* that is not infectious, and exists only within a host cell, being very fragile, often branched, and appearing larger and less dense than the EB.

"Infection-specific" refers to a protein that shows enhanced expression in the RB form of *Chlamydia* compared to the EB form. Infection-specific proteins are not necessarily absent from the EB form, but they are significantly more common in the RB form than in the EB form.

10 "infection-specific antibody" is an antibody that binds specifically to an infection-specific protein.

"Biological sample" refers to any sample of biological origin including, but not limited to a blood sample, a plasma sample, a mucosal smear or a tissue sample.

15 "Isolated" An isolated nucleic acid has been substantially separated or purified away from other nucleic acid sequences in the cell of the organism in which the nucleic acid naturally occurs, i.e., other chromosomal and extrachromosomal DNA and RNA. The term "isolated" thus encompasses nucleic acids purified by standard nucleic acid purification methods. The term also embraces nucleic acids prepared by recombinant expression in a host cell as well as chemically synthesized nucleic acids.

20 "Probes" and "primers." Nucleic acid probes and primers may readily be prepared based on the nucleic acid sequences provided by this invention. A "probe" comprises an isolated nucleic acid attached to a detectable label or reporter molecule. Typical labels include radioactive isotopes, ligands, chemiluminescent agents, and enzymes.

25 "Primers" are short nucleic acids, typically DNA oligonucleotides 15 nucleotides or more in length, which are annealed to a complementary target DNA strand by nucleic acid hybridization to form a hybrid between the primer and the target DNA strand, then extended along the target DNA strand by a DNA polymerase enzyme. Primer pairs can be used for amplification of a nucleic acid sequence, e.g., by the polymerase chain reaction (PCR) or other nucleic-acid amplification methods known in the art.

30 Probes and primers as used in the present invention typically comprise at least 15 nucleotides of the nucleic acid sequences that are shown to encode infection-specific proteins. In order to enhance specificity, longer probes and primers may also be employed, such as probes and primers that comprise at least 20, 30 or 40 consecutive nucleotides of the disclosed nucleic acid sequences.

35 Methods for preparing and using probes and primers are well known in the art and are described in, for example Sambrook et al. (1989); Ausubel et al., (1987); and Innis et al., (1990). PCR primer pairs can be derived from a known sequence, for example, by using computer

programs intended for that purpose such as Primer (Version 0.5, 1991, Whitehead Institute for Biomedical Research, Cambridge, MA).

- “Conservative amino acid substitutions” are those substitutions that, least interfere with the properties of the original protein, i.e., the structure and especially the function of the protein is conserved and not significantly changed by such substitutions. The table below shows amino acids which may be substituted for an original amino acid in a protein and which are regarded as conservative substitutions.

Original Residue	Conservative Substitution
Ala	Ser
Arg	Lys
Asn	gln, his
Asp	Glu
Cys	Ser
Gln	Asn
Glu	Asp
Gly	Pro
His	asn, gln
Ile	leu, val
Leu	ile, val
Lys	arg, gln, glu
Met	leu, ile
Phe	met, leu, tyr
Ser	Thr
Thr	Ser
Trp	Tyr
Tyr	trp, phe
Val	ile, leu

- Conservative substitutions generally maintain (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain.

- The substitutions which in general are expected to produce the greatest changes in protein properties will be non-conservative, for instance changes in which (a) a hydrophilic residue, e.g., seryl or threonyl, is substituted for (or by) a hydrophobic residue, e.g., leucyl, isoleucyl, phenylalanyl, valyl or alanyl; (b) a cysteine or proline is substituted for (or by) any other residue; (c) a residue having an electropositive side chain, e.g., lysyl, arginyl, or histadyl, is substituted for (or by) an electronegative residue, e.g., glutamyl or aspartyl; or (d) a residue having a bulky side chain, e.g., phenylalanine, is substituted for (or by) one not having a side chain, e.g., glycine.

- “Sequence identity” The similarity between two nucleic acid sequences, or two amino acid sequences is expressed in terms of the level of sequence identity shared between the sequences. Sequence identity is typically expressed in terms of percentage identity; the higher the percentage, the more similar the two sequences are. Variants of naturally occurring infection-specific peptides useful in the present invention are typically characterized by possession of at least 50% sequence identity counted over the full length alignment with the amino acid sequence of a

naturally occurring infection-specific peptide when aligned using BLAST 2.0.1 (Altschul et al., 1997). For comparisons of amino acid sequences of greater than about 30 amino acids, the BLAST 2 analysis is employed using the blastp program set to default parameters (open gap = 11, extension gap = 1 penalty, gap x dropoff = 50, expect = 10, word size = 3, filter on), and using

5 the default BLOSUM62 matrix (gap existence cost = 11, per residue gap cost = 1, lambda ratio = 0.85). When aligning short peptides (fewer than around 30 amino acids), the alignment should be performed using the Blast 2 sequences function, employing the PAM30 matrix (gap existence cost = 9, per residue gap cost = 1, lambda ratio = 0.87). Proteins with even greater similarity to the reference sequences will show increasing percentage identities when assessed by this method,

10 such as at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, or at least 95% sequence identity. The NCBI Basic Local Alignment Search Tool (BLAST) (Altschul et al., 1990) is available from several sources, including the National Center for Biotechnology Information (NCBI, Bethesda, MD) and on the Internet, for use in connection with the sequence analysis programs blastp, blastn, blastx, tblastn and tblastx. It can be accessed at

15 <http://www.ncbi.nlm.nih.gov/BLAST/>. A description of how to determine sequence identity using this program is available at http://www.ncbi.nlm.nih.gov/BLAST/blast_help.html.

Similarly, when comparing nucleotides, blastn may be used with default settings (rewards for match = 1, penalty for mismatch = -2, open gap = 5, extension gap = 2 penalty, gap x dropoff = 50, expect = 10, word size = 11, filter on), with the default BLOSUM62 matrix (as

20 above). Variants of naturally occurring infection-specific nucleic acid sequences useful in the present invention are typically characterized by possession of at least 50% sequence identity counted over the full length alignment with the nucleic acid sequence of a naturally occurring infection-specific ORF when aligned using BLAST 2.0.1. Useful nucleic acids may show even greater percentage identity, and may, for example, possess at least 55%, at least 65%, at least

25 75%, at least 80%, at least 85%, at least 90%, or at least 95% sequence identity naturally occurring infection-specific ORF.

"Operably linked" A first nucleic acid sequence is "operably" linked with a second nucleic acid sequence when the first nucleic acid sequence is placed in a functional relationship with the nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence

30 if the promoter affects the transcription or expression of the coding sequence. Generally, operably linked DNA sequences are contiguous and, where necessary to join two protein coding regions, in the same reading frame.

"Recombinant" A recombinant nucleic acid is one that has a sequence that is not naturally occurring or has a sequence that is made by an artificial combination of two otherwise separated

35 segments of sequence. This artificial combination is often accomplished by chemical synthesis or, more commonly, by the artificial manipulation of isolated segments of nucleic acids, e.g., by genetic engineering techniques.

"Stringent Conditions" Stringent conditions, in the context of nucleic acid hybridization, are sequence-dependent and are different under different environmental parameters. Generally, stringent conditions are selected to be about 5 degrees to 20 degrees lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. Conditions for nucleic acid hybridization and calculation of stringencies can be found in Sambrook et al. (1989), pages 9.49-9.55. Typical high stringency hybridization conditions (using radiolabeled probes to hybridize to nucleic acids immobilized on a nitrocellulose filter) may include, for example, wash conditions of 0.1 X SSC, 0.5% SDS at a wash temperature of 68°C.

When referring to a probe or primer, the term "specific for (a target sequence)" indicates that the probe or primer hybridizes under high-stringency conditions substantially only to the target sequence in a given sample comprising the target sequence.

"Purified" A purified peptide is a peptide that has been extracted from the cellular environment and separated from substantially all other cellular peptides. As used herein, the term peptide includes peptides, polypeptides and proteins. In certain embodiments, a purified peptide is a preparation in which the subject peptide comprises 50% or more of the protein content of the preparation. For certain uses, such as vaccine preparations, even greater purity may be preferable.

"Immunostimulatory peptide" as used herein refers to a peptide that is capable of stimulating a humoral or antibody-mediated immune response when inoculated into an animal.

"Vaccine" A vaccine is a composition containing at least one immunostimulatory peptide which may be inoculated into an animal with the intention of producing a protective immunological reaction against a certain antigen. The antigen to be protected against may be, for instance, an infectio-specific antigen of *Chlamydia*.

B. ISOLATION OF INFECTION SPECIFIC CHLAMYDIA POLYPEPTIDES AND IDENTIFICATION OF GENES ENCODING THESE POLYPEPTIDES

1. ISOLATION OF IncA, IncB AND IncC

Bacterial strains. *Chlamydia* (*C. psittaci* strain GPIC or *C. trachomatis* LGV-434, ser. L2) was cultivated in HeLa 229 cells using standard methods (Caldwell et al., 1981). Purified *Chlamydiae* were obtained using Renografin (E. R. Squibb & Sons, Inc., Princeton, N.J.) density gradient centrifugation. *Escherichia coli* DH5 α (Bethesda Research Laboratories, Inc., Gaithersburg, Md.) was used as the host strain for transformations with recombinant DNA. *E. coli* XL1-Blue MRF' (Stratagene, La Jolla, Calif.) was used as the host strain for infection with lambda ZAPII phage vector. *E. coli* SOLR (Stratagene) was used as the host strain for infection with *in vivo* excised filamentous lambda ZAPII.

Antisera. MBP (Maltose Binding Protein)-Inc fusion proteins were used as antigens for the production of mono-specific antibody reagents in Hartley strain guinea-pigs. The protein was diluted to 100 $\mu\text{g}/\text{ml}^{-1}$ sterile saline and mixed with the Ribí Trivalent Adjuvant (Ribí Immunochem.). The antigen/adjuvant emulsion was administered to anaesthetized guinea-pigs using a procedure provided by the manufacturer. Sera were collected 14 days after secondary and tertiary immunizations. Control antisera were produced by immunizing guinea-pigs with adjuvant alone, or with adjuvant plus purified maltose-binding protein.

Convalescent guinea-pig antisera, antisera against live EBs, and antisera against formalin-fixed EBs were produced using standard methods (Rockey and Rosquist, 1994 and Rocky et al., 1995).

C. psittaci library construction and screening. For the *incB* and *incC* genes, *C. psittaci* strain GPIC DNA was extracted using a genomic DNA extraction kit (Qiagen) with one modification; dithiothreitol (5mM) was added to the suspension buffer to assist EB lysis. DNA was partially digested with *Tsp509I* and ligated to *EcoRI* digested λ -ZAPII phage arms (Stratagene). The ligation was packaged in vitro with Gigapack extracts according to the manufacturer's instructions (Stratagene). Recombinant phage were plated on *E. coli* XL-1 Blue at densities of approximately 10^4 PFU/150-mm (diameter) plate. Following a nine hour incubation to allow development of the plaques, the plates were sequentially overlaid with nitrocellulose disks and the resulting lifts were processed for immunoblotting with convalescent antisera and antisera to fixed EBs. Of approximately 8,000 plaques, 18 had reactivity with the convalescent sera but not sera generated against EBs. One of these was subcloned into pBluescript SK(-) phagmid by *in vitro* excision in the *E. coli* SOLR strain (Stratagene) and designated pBS200-7.

For the *incA* gene, genomic DNA from *C. psittaci* strain GPIC was partially digested with *Sau3A*, size-selected (2-8 kb) by electrophoresis through low-melting-temperature agarose, and blunt-ended with T4 DNA polymerase. This DNA was ligated to an *EcoRI*/*NotI* adapter (Life Technologies), kinased, and ligated to *EcoRI*-digested Lambda ZAP II vector (Stratagene Cloning Systems). Recombinants were packaged (Lambda Gigapack Gold, Stratagene) and used to infect *E. coli* XL1-Blue (Stratagene). Plaques were allowed to develop for 4 h at 37°C. Nitrocellulose filters laden with 10 mM IPTG (US Biochemical Corp.) were placed onto the plaques and incubated for an additional 4 h at 37°C. These filters were removed and placed into a blocking solution consisting of PBS (150 mM NaCl, 10 mM NaPO₄, pH7.2) plus 0.1 % Tween-20 (TPBS) and 2% BSA-TPBS. Filters were incubated for 1 h, rinsed twice in TPBS, and incubated overnight in convalescent-guinea-pig sera diluted 1:100 in BSA-TPBS. After three washes in TPBS, the filters were incubated for 1 h in ¹²⁵I-staphylococcal protein A (New England Nuclear) diluted to approx. 124 nCi/ml⁻¹ in BSA-TPBS. Filters were again washed three times in TPBS and positive plaques were detected by exposure of the dried filters to autoradiography film overnight at room temperature. Positive clones were picked and plaque-purified. pBluescript-SK- plasmids

containing the chlamydial genes of interest were recovered from the purified bacteriophage using ExAssist filamentous bacteriophages (Stratagene).

Identification of antigens recognized by convalescent antisera. Recombinant plaques were identified that showed reactivity with convalescent (anti-RB) antisera, but not with anti-EB serum. The purified recombinant phage were converted into pBluescriptII SK plasmid by *in vivo* excision and recircularization and these recombinant DNAs were used to transform *E. coli*. SDS-PAGE and immunoblot analysis of lysates of these recombinant *E. coli* showed that each expressed one or more proteins that reacted with convalescent antisera but not with the EB serum.

DNA Cloning and fusion protein production. The plasmid pJC2 contains a 5.0 kb *EcoRI* GPIC genomic fragment cloned into the pZero2.1 vector (Invitrogen). To construct pJC2, the *incC* ORF sequence was ³²P-radiolabeled using random priming (Gibco-BRL) and used to probe *EcoRI* cut GPIC genomic DNA fragments separated by agarose gel electrophoresis. Fragments in the size range of the positive signal were excised from the gel and purified by Gene-Clean (Bio101). The gel-purified fragments were used in a ligation along with *EcoRI*-digested pZero2.1. Kanamycin resistant colonies were screened by colony hybridization with radiolabeled *incC*.

MBP fusions of the five ORFs present in pJC2 were produced using the pMAL-C2 vector (New England Biolabs). The reading frame of *incC*, with the exception of the first four codons, was amplified using *Pwo* polymerase (Boehringer Mannheim) and pBS200-7 as the template. The upstream and downstream oligonucleotides for this amplification were

5'-AGAACCGATTTAACTCCAGGCG-3' (SEQ ID NO: 19) and

5'-GCGCGGATCCTTAATGTCCGGTAGGCCTAG-3' (SEQ ID NO: 20), respectively.

The vector was digested with *XmnI* and *BamHI*, and the amplification product was digested with *BamHI*. Ligation of these products resulted in an in-frame fusion between the *malE* gene in the vector and the *incC* reading frame from pBS200-7. The stop codon for this construction is provided by the insert. Following ligation, the products were transformed into *E. coli* strain HD50. The resulting fusion protein (MBP/IncC) was overexpressed and purified by maltose affinity chromatography using an amylose resin supplied by New England Biolabs.

The same approach was used for production of the MBP/IncB fusion protein. The sequence encoding the N-terminal 101 amino acids of the IncB ORF was PCR amplified using the oligonucleotides

5'-ATGTCAACAACACCAGCATCTTC-3' (SEQ ID NO: 21) and

5'-GCGCGGATCCTTAATTAGTGCCTTCTGGATTAGG-3' (SEQ ID NO: 22).

The purified MBP/IncB and MBP/IncC fusion proteins were used as antigen for the production of monospecific antibody in Hartley strain guinea-pigs by standard methods (Rockey et al., 1995). Inserts in each construct were confirmed by DNA sequencing.

For IncA, a maltose-binding protein/IncA fusion protein was produced using the pMAL-C2 vector system from New England Biolabs. The reading frame of *incA* shown in Fig.1, with the exception of the initiator ATG, the *incA* ORF was amplified using Vent DNA polymerase (New England Biolabs) and plasmid pGP17 as template. The upstream and downstream oligo-

5 nucleotides for this amplification were

5'-CGCAGTACTGTATCCACAGACAAC-3' (SEQ ID NO: 23) and

5'-GTCGGATCCGAGAACTCTCCATGCC-3' (SEQ ID NO: 24), respectively. The vector was digested with *XmnI* and *Bam*HI, and the amplification product was digested with *ScaI* and *Bam*HI. Ligation of these products resulted in an in-frame fusion between the *malE* gene in the vector and the *incA* reading frame from pGP17. The stop codon for this construction is provided by the insert. Following ligation, the products were transformed into *E. coli* strain DH5 α . The resulting fusion protein (MBP/IncA) was overexpressed and purified by maltose affinity chromatography using amylose resin (New England Biolabs).

10

MBP/IncA was used as antigen for the production of mono-specific antibody reagents in Hartley strain guinea-pigs.

15

DNA sequencing and sequence analysis. The pBS200-7 and pJC2 genomic clones as well as the MBP fusions were sequenced with the *Taq* DyeDeoxy Terminator Cycle Sequencing Kit (Perkin Elmer/Applied Biosystems Division). Several internal primers were designed to sequence further into the cloned inserts. Sequence assembly was performed using AssemblyLIGN software and sequence analysis was performed with MacVector software (International Biotechnologies Incorporated). Hydrophilicity profiles were determined using the Kyte-Doolittle scale (Kyte and Doolittle, 1982) with a window of 7. Deduced amino acid sequences were compared with the database using the BLAST program (on default settings) available from the National Center for Biotechnology Information on the world wide web. The entire nucleotide sequence of the pJC2 insert was deposited in the GenBank/EMBL Nucleotide Sequence Data Library, under accession number AF017105.

20

25

For *incA*, nucleotide sequencing was conducted using the Sequences system (US Biochemical) with the M13 forward and reverse primers, and internal primers synthesized on a Milligen/Bioscience Cyclone Plus DNA synthesizer. Computer analyses were conducted using the MacVector Sequence Analysis Software (International Biotechnologies Incorporated). Hydrophilicity profiles were determined using the Kyte-Doolittle scale (Kyte and Doolittle, 1982) with a window of 7. Secondary-structure predictions were generated using a combination of the Chou-Fasman and Robson-Garnier methods (Robson and Suzuki, 1976; Chou and Fasman, 1978). Deduced amino acid sequences were compared with those in the EMBL and GenBank databases using the BLASTP program available from the National Center for Biotechnology Information.

30

35

Electrophoresis and immunoblotting. Polyacrylamide gel electrophoresis (PAGE) was conducted using standard methods (Rockey and Rosquist, 1994). Immunoblotting was performed using standard methods (Rockey et al., 1995).

Immunofluorescence studies. *Chlamydiae* grown in HeLa cells on sterile glass coverslips were fixed for microscopy one of two ways. Cells were either incubated in methanol for 5 minutes, or in the combination fixative periodate-lysine-paraformaldehyde (PLP) for three hours at room temperature followed by permeabilization with 0.05% saponin (Brown and Farquhar, 1989). Immunostaining of the fixed coverslips was performed according to standard methods (Rockey et al., 1995) and visualized under a Nikon Microphot FXA microscope using the 63x objective and oil immersion.

RT-PCR analysis. RNA for RT-PCR analysis was extracted from approximately 2×10^4 *C. psittaci*-infected cells. A Qiagen column was used for extraction and purification according to the manufacturer's instructions (Qiagen). RQ1 RNase DNase (Promega) was used to ensure removal of contaminating genomic DNA. cDNA was prepared by incubating 1.5 μ g of RNA, 2.5 μ M of the reverse oligonucleotide primer, and AMV reverse transcriptase (Promega) for 1 hour at 42°C in sodium pyrophosphate buffer, according to the manufacturer's instructions. PCR reactions were carried out using 1 μ l of the cDNA reaction, 1.25 μ M of each oligonucleotide primer, and *Pwo* polymerase (Boehringer Mannheim). Each RT-PCR reaction was accompanied by a positive control reaction that utilized the same primer set and 10 ng of *C. psittaci* genomic DNA, and a negative control reaction in which 1 μ l of the same RNA preparation was used as template in the PCR reaction. A control primer set located within the *incC* gene was also used as an RT-PCR control.

Identification of *incA*, *incB* and *incC* genes of *C. trachomatis*. The nucleotide sequence information obtained for the *incA*, *incB* and *incC* of *C. psittaci* (above) was used, with standard methods, to identify the *inc* gene orthologues of *C. trachomatis*. Probes were made that corresponded to the 3' and 5' ends of the *C. psittaci inc* open reading frames. Standard PCR amplification (as above) was used, with the *C. trachomatis* genome as a template, to amplify the corresponding *C. trachomatis* nucleotide sequence. The amplified DNA was then sequenced, using standard methods.

2. ISOLATION OF p242, TroA AND TroB

Bacterial strains. *C. trachomatis* LGV-434, serotype L2, was cultivated in HeLa 229 cells using standard methods (Caldwell et al., 1981). Purified *chlamydiae* were obtained using Renografin (E. R. Squibb & Sons, Inc., Princeton, N.J.) density gradient centrifugation (Hackstadt et al., 1992). *Escherichia coli* DH5 α (Bethesda Research Laboratories, Inc., Gaithersburg, Md.) was used as the host strain for transformations with recombinant DNA. *E. coli* XL1-Blue MRF' (Stratagene, La Jolla, Calif.) was used as the host strain for infection with lambda ZAPII phage

vector. *E. coli* SOLR (Stratagene) was used as the host strain for infection with *in vivo* excised filamentous lambda ZAPII.

Antisera. Two Cynomolgus monkeys (*Macaca fascicularis*) were anaesthetized and infected urethrally with *C. trachomatis* EBs. Each monkey was infected twice and allowed to
5 recover between infections. Symptoms of infection were monitored over time. Antisera from infected monkeys were tested for reactivity to *Chlamydia* by ELISA (Su et al., 1990).

Sera were collected every two weeks and anti-chlamydial titers were determined. These animals showed mild clinical signs of disease which cleared spontaneously. A second challenge was then administered. Sera were collected from these animals and used to probe a *C. trachomatis*
10 expression library as discussed below. As a control, Guinea Pigs were immunized with killed *C. trachomatis* of the EB form. Sera from these animals were obtained and also used to probe the *C. trachomatis* expression library.

***C. trachomatis* library construction and immunoscreening.** A *C. trachomatis* genomic library was constructed with the lambda ZAPII vector as described above for *C. psittaci*.

15 Approximately 15,000 plaques were plated, transferred to nitrocellulose filters (Schleicher and Schuell, Keene, N.H.) in duplicate, and probed with the monkey convalescent antiserum and with Guinea Pig serum against killed EBs (Bannantine et al., 1998). Plaques that reacted only with the monkey convalescent antisera were selected for further study.

Identification of antigens recognized by convalescent antisera. Four positive
20 recombinant plaques were identified that showed reactivity with convalescent antisera but not with anti-EB serum. The purified recombinant phage were converted into pBluescriptII SK plasmid by *in vivo* excision and recircularization and these recombinant DNAs (pCt1, pCt2, pCt3 and pCt4) were used to transform *E. coli*. SDS-PAGE and immunoblot analysis of lysates of these recombinant *E. coli* showed that each expressed one or more proteins that reacted with
25 convalescent (anti-RB) antisera but not with the anti-EB antiserum. Two of the recombinants clones, pCt2 and pCt3, expressed an identical 19.9 kDa protein (p242). The pCt4 recombinant expressed two different proteins of approximately 32 kDa each that are strongly recognized by convalescent antisera (TroA and TroB).

30 C. SEQUENCE ANALYSIS

Sequence analysis of pCt1, 2, and 3 revealed overlapping inserts with only one open reading frame (ORF) common in all three. This ORF encodes an approximately 19.9 kDa protein (p242) that shows no similarity to other known proteins. The nucleotide sequence encoding *C. trachomatis* p242, and the amino acid sequence of the protein are shown in SEQ ID NOS:1 and 2,
35 respectively.

The insert in pCt4 contains two complete ORFs which code for two proteins, each of approximately 32kDa (TroA and TroB) that show some homology with proteins from *Treponema*

pallidum. The nucleotide sequences encoding the 32 kDa proteins (TroA and TroB) and the amino acid sequences of these proteins are shown in SEQ ID NOS: 3, 4, 5, and 6.

D. EMBODIMENTS OF THE INVENTION

5 The present invention includes the nucleotide and amino acid sequences for certain infection-specific proteins from *Chlamydia*. These proteins are p242, TroA, and TroB from *C. trachomatis*, and the IncB, and IncC proteins from *C. psittaci*. The scope of the invention includes fragments of these proteins that may be used in a vaccine preparation or that may be used in a method of detecting *Chlamydia* antibodies. Such fragments may be, for example, 5, 10, 15, 10 20, 25, or 30 contiguous amino acids in length, or may even encompass the entire protein.

The present invention also encompasses the use of infection-specific proteins of *Chlamydia*, and the use of nucleotides encoding such proteins. Infection-specific proteins include the IncA, IncB and IncC proteins of *C. psittaci*, the IncA, IncB and IncC proteins of *C. trachomatis*, and the TroA, TroB, and p242 proteins of *C. trachomatis*. The inventors have shown 15 that these proteins are infection-specific by using immunological techniques such as immunofluorescence microscopy and immunoblotting.

The present invention includes a vaccine against chlamydial infections comprising infection-specific proteins or fragments of these proteins or proteins that are homologous or show substantial sequence similarity to these proteins. In one embodiment, one or more purified 20 infection-specific proteins may be mixed with a pharmaceutically acceptable excipient to produce a vaccine that stimulates a protective immunological response in an animal. In one embodiment the vaccine may be administered intra-muscularly or sub-cutaneously or intravenously. In another embodiment, the vaccine may be administered by inoculation into or onto the mucous membranes of the subject animal. For example, the vaccine may be administered urethrally or genitally as a liquid or in the form of a pessary. In another embodiment, it may be administered to the mucosa 25 of the lungs as a spray or vapor suspension.

Since at least three amino acids are required to produce an antigenic epitope, the vaccine should comprise at least three consecutive amino acids, preferably at least five consecutive amino acids, and may comprise at least 10, 15, 25, 30, 40, or 45 consecutive amino acids of the 30 infection-specific proteins as shown in SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, and 18.

The vaccine of the invention may be used to inoculate potential animal targets of any of the chlamydial diseases including those caused by *C. psittaci*, *C. trachomatis*, *C. pneumoniae* or *C. pecorum*. Indeed the vaccine of the invention may be used to inoculate animals against any disease that shows immunological cross-protection as a result of exposure to infection-specific 35 *Chlamydia* antigen.

Vaccines of the present invention can include effective amounts of immunological adjuvants known to enhance an immune response (e.g., alum). The protein or polypeptide is present in the vaccine in an amount sufficient to induce a protective immune response whether

through humoral or cell mediated pathways or through both. Such a response protects the immunized animal against chlamydial infections specifically by raising an immune response against the Reticulate Body form of *Chlamydia*. Protective antibodies may be elicited by a series of two or three doses of the antigenic vaccine given about two weeks apart.

5 The present invention also teaches a method of making a vaccine against chlamydial infections. The method of making the vaccine comprises providing a pure (or substantially pure) infection-specific chlamydial peptide or portion thereof, and mixing the peptide with a pharmacologically acceptable excipient or adjuvant. Adjuvants may include commonly used compounds such as alum. Additionally, the vaccines may be formulated using a peptide according
10 to the present invention together with a pharmaceutically acceptable excipient such as water, saline, dextrose and glycerol. The vaccines may also include auxiliary substances such as emulsifying agents and pH buffers. Doses of the vaccine administered will vary depending on the antigenicity of the particular peptide or peptide combination employed in the vaccine and characteristics of the animal or human patient to be vaccinated.

15 The infection-specific vaccine of the invention is directed towards not only *C. psittaci*, but against all forms of *Chlamydia* including *C. pneumoniae*, *C. trachomatis* and *C. pecorum*, and the vaccine may comprise not just peptides derived from *C. psittaci*, but also orthologous peptides and fragments of such orthologous peptides from other species of *Chlamydia* and peptides that are substantially similar to such peptides.

20 The present invention also teaches a method of vaccination comprising administering a vaccine formulated as described above to an animal either intravenously, intramuscularly, subcutaneously, by inhalation of a vapor or mist, or by inoculation in the form of a liquid, spray, ointment, pessary or pill into or onto the mucous membranes of the mouth, nose, lungs or urogenital tract or colon.

25 The methods of the invention may be practiced equally with human or non-human animal subjects.

 The present invention also teaches a method of detecting *Chlamydia* infection-specific proteins produced by the Reticulate Body form of the organism. In this embodiment, antibodies raised to the infection-specific proteins are used in an immunological assay such as an Enzyme
30 Linked Immunosorbant Assay or Biotin-Avidin assay or a radioimmunoassay or any other assay wherein specific antibodies are used to recognize a specific protein. Such assays may be used to detect both the quantity of proteins present and also the specificity of binding of such proteins. In such an assay, antibodies have attached to them, usually at the *Fc* portion, a detectable label, such as an enzyme, fluorescent marker, a radioactive marker or a Biotin-Avidin system marker that
35 allows detection. A biological sample is provided from an animal that has been putatively exposed to *Chlamydia*. Such a sample may be, for example, whole blood, serum, tissue, saliva or a mucosal secretion. The sample is then contacted with the labeled antibody and specific binding, if any, is detected. Other methods of using infection-specific antibodies to detect infection-specific

antigens that are present in cells or tissues include immunofluorescence, indirect-immunofluorescence and immunohistochemistry. In immunofluorescence, a fluorescent dye is bound directly to the antibody. In indirect-immunofluorescence, the dye is bound to an anti-immunoglobulin. Specific binding occurs between antigen and bound antibody is detected by virtue of fluorescent emissions from the dye moiety. This technique would be particularly useful, for instance, for detection of *Chlamydia* antigen present on a urogenital mucosal smear.

Other techniques, such as competitive inhibition assays may also be used to assay for antigen, and one of ordinary skill in the art will readily appreciate that the precise methods disclosed may be modified or varied without departing from the subject or spirit of the invention taught herein.

The present invention also teaches a method of detection of *Chlamydia* infection-specific antibodies made against the Reticulate Body. In this embodiment a sample is provided from an animal putatively exposed to *Chlamydia* to determine whether the sample contains infection-specific antibodies. Such a sample may be, for example, whole blood, serum, tissue, saliva or a mucosal secretion. This sample is contacted with infection-specific antigens such that the amount and specificity of binding of the antibody may be measured by its binding to a specific antigen. Many techniques are commonly known in the art for the detection and quantification of antigen. Most commonly, the purified antigen will be bound to a substrate, the antibody of the sample will bind via its *Fab* portion to this antigen, the substrate will then be washed and a second, labeled antibody will then be added which will bind to the *Fc* portion of the antibody that is the subject of the assay. The second, labeled antibody will be species specific, i.e., if the serum is from a human, the second, labeled antibody will be anti-human-IgG antibody. The specimen will then be washed and the amount of the second, labeled antibody that has been bound will be detected and quantified by standard methods.

The present invention also teaches a method of treating a *Chlamydial* infection by directing a therapeutic agent against a specific target, such as: (i) an infection-specific protein of *Chlamydia*, (ii) a gene that encodes an infection-specific protein of *Chlamydia* and (iii) an RNA transcript that encodes an infection-specific protein of *Chlamydia*, wherein said therapeutic agent interacts with said target to affect a reduction in pathology.

For example, the present invention teaches a method of treating chlamydial infection wherein antisense technology is used to prevent the expression of infection-specific genes, thereby preventing the pathologies associated these proteins and preventing reproduction of the RB phase of *Chlamydia*. In this embodiment, RNA molecules complementary to transcripts of infection specific genes are introduced into the host cells that contain *Chlamydia*, and by binding to the mRNA transcripts of the infection-specific genes, prevent translation and therefore expression of the infection-specific proteins that are associated with pathogenesis.

The invention may be practiced to produce a vaccine against any species of *Chlamydia*, including *C. psittaci*, *C. pecorum*, *C. trachomatis* and *C. pneumoniae*.

The following examples illustrate various embodiments of the invention.

EXAMPLE 1: Homologous Sequences

The DNA and protein sequences discussed herein are shown in SEQ ID NOS:1-18.

5 These sequences refer to infection-specific proteins and to the DNA sequences that encode these proteins. Although these sequences are from *C. psittaci* and *C. trachomatis*, it would be equally possible to substitute in the present invention, the orthologs of these sequences from other *Chlamydia* species such as *C. pecorum* and *C. pneumoniae*.

10 Such orthologous sequences may be obtained from the appropriate organisms by isolation of the genome of the organism, digestion with restriction enzymes, separation of restriction fragments by electrophoresis and purification of these fragments and selection of fragments of appropriate size. Identity of the fragments can be confirmed by dot-blot and by standard DNA sequencing techniques. The orthologous sequences in different *Chlamydia* species may also be found by selection of appropriate PCR primers (selected from appropriate regions flanking the

15 *Chlamydia* gene of interest), and the use of these primers in a PCR reaction, using the genome of the particular species of *Chlamydia* of interest as a template, to amplify the ortholog of interest. Such PCR primers would be selected from the flanking regions to allow specific amplification of the target gene. The fragments so obtained could then be run on a gel to check size and sequenced and compared against the known sequences to determine sequence identity.

20 The degree of sequence identity between the infection-specific genes of *C. psittaci* or *C. trachomatis* and their orthologs from *C. pecorum* and *C. pneumoniae*, may be determined by comparing sequences using the National Center for Biotechnology Information (NCBI) Basic Local Alignment Search Tool (BLAST) as described herein.

Orthologues of interest infection-specific proteins are characterized by possession of at

25 least 50% or greater sequence identity counted over the full length alignment with one of the disclosed amino acid sequences of the *C. psittaci* or *C. trachomatis* infection-specific proteins using gapped blastp set to default parameters (described herein).

EXAMPLE 2: Heterologous Expression of Infection-Specific Antigens

30 Methods for expressing large amounts of protein from a cloned gene introduced into *Escherichia coli* (*E. coli*) may be utilized for the purification of the *Chlamydia* peptides. Methods and plasmid vectors for producing fusion proteins and intact native proteins in bacteria are well known and are described in Sambrook et al. (1989). Such fusion proteins may be made in large amounts, are relatively simple to purify, and can be used to produce antibodies. Native proteins

35 can be produced in bacteria by placing a strong, regulated promoter and an efficient ribosome binding site upstream of the cloned gene. If low levels of protein are produced, additional steps

may be taken to increase protein production; if high levels of protein are produced, purification is relatively easy.

Often, proteins expressed at high levels are found in insoluble inclusion bodies. Methods for extracting proteins from these aggregates are described in chapter 17 of Sambrook et al.

5 (1989). Vector systems suitable for the expression of *lacZ* fusion genes include the pUC series of vectors (Ruther et al. (1983)), pEX1-3 (Stanley and Luzio (1984)) and pMR100 (Gray et al. (1982)). Vectors suitable for the production of intact native proteins include pKC30 (Shimatake and Rosenberg (1981)), pKK177-3 (Amann and Brosius (1985)) and pET-3 (Studiar and Moffatt (1986)).

10 Fusion proteins may be isolated from protein gels, lyophilized, ground into a powder and used as antigen preparations.

Mammalian or other eukaryotic host cells, such as those of yeast, filamentous fungi, plant, insect, amphibian or avian species, may also be used for protein expression, as is well known in the art. Examples of commonly used mammalian host cell lines are VERO and HeLa
15 cells, Chinese hamster ovary (CHO) cells, and WI38, BHK, and COS cell lines, although it will be appreciated by the skilled practitioner that other prokaryotic and eukaryotic cells and cell lines may be appropriate for a variety of purposes, e.g., to provide higher expression, post-translational modification, desirable glycosylation patterns, or other features.

Additionally, peptides, particularly shorter peptides, may be chemically synthesized,
20 avoiding the need for purification from cells or culture media. It is known that peptides as short as 3 amino acids can act as an antigenic determinant and stimulate an immune response. Such peptides may be administered as vaccines in ISCOMs (Immune Stimulatory Complexes) as described by Janeway & Travers, Immunobiology: The Immune System In Health and Disease, 13.21 (Garland Publishing, Inc. New York, 1997). Accordingly, one aspect of the present
25 invention includes small peptides encoded by the nucleic acid molecules disclosed herein. Such peptides include at least 5, and may be at least 10, 15, 20, 25, or 30 or more contiguous amino acids of the polypeptide sequences described herein.

30 **EXAMPLE 3: Production of Antibodies Specific for Infection-Specific Antigens**

Antibody against infection-specific antigen is encompassed by the present invention, particularly for the detection of *Chlamydia* infection-specific antigen. Such antibody may be produced by inoculation of an animal such as a guinea-pig or a monkey with infection-specific antigen produced as described above. Such antigen may be a polypeptide as disclosed herein, such
35 as a complete or partial polypeptide from *C. psittaci*, *C. trachomatis*, *C. pneumoniae* or *C. pecorum*. As discussed above, any molecule that can elicit a specific, protective immune response

may be used as a vaccine, but since a minimum of three amino acids are required to do this, a vaccine should comprise at least three amino acids.

The peptide for use in the vaccine of the invention may be naturally derived or may be synthetic such as those synthesized on a commercially available peptide synthesizer. The peptide
5 may also comprise a complete or partial peptide derived from the *C. pneumoniae* or *C. pecorum* infection-specific orthologs of the *C. trachomatis* or *C. psittaci* proteins as set out herein.

In one method of production, a polyclonal antibody is produced by providing a purified peptide which is diluted to 100 micrograms per milliliter in sterile saline and mixed with RiBi Trivalent Adjuvant (RiBi Immunochem Inc). The antigen/adjuvant emulsion is then administered
10 to an anaesthetized guinea pig using a procedure as provided by the manufacturer. Serum is collected 14 days after secondary and tertiary immunizations.

Monoclonal antibody to epitopes of the *Chlamydia* peptides identified and isolated as described can be prepared from murine hybridomas according to the classical method of Kohler and Milstein (1975) or derivative methods thereof. Briefly, a mouse is repetitively inoculated with
15 a few micrograms of the selected purified protein over a period of a few weeks. The mouse is then sacrificed, and the antibody-producing cells of the spleen isolated. The spleen cells are fused by means of polyethylene glycol with mouse myeloma cells, and the excess unfused cells destroyed by growth of the system on selective media comprising aminopterin, e.g., Hypoxanthene, Aminopterin and Thymidine (HAT) medium. The successfully fused cells are diluted and aliquots
20 of the dilution placed in wells of a microtiter plate where growth of the culture is continued. Antibody-producing clones are identified by detection of antibody in the supernatant fluid of the wells by immunoassay procedures, such as ELISA, as originally described by Engvall (1980), and derivative methods thereof. Selected positive clones can be expanded and their monoclonal antibody product harvested for use. Detailed procedures for monoclonal antibody production are
25 described in Harlow and Lane (1988).

An alternative approach to raising antibodies against the *Chlamydia* peptides is to use synthetic peptides synthesized on a commercially available peptide synthesizer based upon the amino acid sequence of the peptides predicted from nucleotide sequence data.

In another embodiment of the present invention, monoclonal antibodies that recognize a
30 specific *Chlamydia* peptide are produced. Optimally, monoclonal antibodies will be specific to each peptide, i.e., such antibodies recognize and bind one *Chlamydia* peptide and do not substantially recognize or bind to other proteins, including those found in uninfected human cells.

The determination that an antibody specifically detects a particular *Chlamydia* peptide is made by any one of a number of standard immunoassay methods; for instance, the western blotting
35 technique (Sambrook et al., 1989). To determine that a given antibody preparation (for instance from a guinea pig) specifically detects one *Chlamydia* peptide by western blotting, total cellular protein is extracted from a sample of blood from an unexposed subject and from a sample of blood from an exposed subject. As a positive control, total cellular protein is also extracted from

Chlamydia cells grown *in vitro*. These protein preparations are then electrophoresed on a sodium dodecyl sulfate-polyacrylamide gel. Thereafter, the proteins are transferred to a membrane (for example, nitrocellulose) by western blotting, and the antibody preparation is incubated with the membrane. After washing the membrane to remove non-specifically bound antibodies, the
5 presence of specifically bound antibodies is detected by the use of an anti-guinea pig antibody conjugated to an enzyme such as alkaline phosphatase; application of the substrate 5-bromo-4-chloro-3-indolyl phosphate/nitro blue tetrazolium results in the production of a dense blue compound by immuno-localized alkaline phosphatase. Antibodies which specifically detect the *Chlamydia* protein will, by this technique, be shown to bind to the *Chlamydia*-extracted sample at
10 a particular protein band (which will be localized at a given position on the gel determined by its molecular weight) and to the proteins extracted from the blood of the exposed subject. No significant binding will be detected to proteins from the unexposed subject.

**EXAMPLE 4: Use of Infection-Specific Sequences
and their Corresponding Peptides and
15 Antibodies in Diagnostic Assays**

Another aspect of the present invention is a method for detecting the presence of anti-*Chlamydia* antibodies that react with infection-specific *Chlamydia* proteins, *Chlamydia* peptides and *Chlamydia* nucleic acid sequences in biological samples. These methods include detection of antigen and antibody by ELISA and similar techniques, the detection of proteins in a tissue sample
20 by immunofluorescence and related techniques and the detection of specific DNA sequences by specific hybridization and amplification.

One aspect of the invention is an ELISA that detects anti-*Chlamydia* antibodies in a medical specimen. An immunostimulatory infection-specific *Chlamydia* peptide of the present invention is employed as an antigen and is preferably bound to a solid matrix such as a crosslinked dextran such as SEPHADEX (Pharmacia, Piscataway, NJ), agarose, polystyrene, or the wells of a
25 microtiter plate. The polypeptide is admixed with the specimen, such as blood, and the admixture is incubated for a sufficient time to allow antibodies present in the sample to immunoreact with the polypeptide. The presence of the positive immunoreaction is then determined using an ELISA assay, usually involving the use of an enzyme linked to an anti-immunoglobulin that catalyzes the
30 conversion of a chromogenic substrate.

In one embodiment, the solid support to which the polypeptide is attached is the wall of a microtiter assay plate. After attachment of the polypeptide, any nonspecific binding sites on the microtiter well walls are blocked with a protein such as bovine serum albumin. Excess bovine serum albumin is removed by rinsing and the medical specimen is admixed with the polypeptide in
35 the microtiter wells. After a sufficient incubation time, the microtiter wells are rinsed to remove excess sample and then a solution of a second antibody, capable of detecting human antibodies is added to the wells. This second antibody is typically linked to an enzyme such as peroxidase,

alkaline phosphatase or glucose oxidase. For example, the second antibody may be a peroxidase-labeled goat anti-human antibody. After further incubation, excess amounts of the second antibody are removed by rinsing and a solution containing a substrate for the enzyme label (such as hydrogen peroxide for the peroxidase enzyme) and a color-forming dye precursor, such as o-phenylenediamine is added. The combination of *Chlamydia* peptide (bound to the wall of the well), the human anti-*Chlamydia* antibodies (from the specimen), the enzyme-conjugated anti-human antibody and the color substrate will produce a color that can be read using an instrument that determines optical density, such as a spectrophotometer. These readings can be compared to a negative control such as a sample known to be free of anti-*Chlamydia* antibodies. Positive readings indicate the presence of anti-*Chlamydia* antibodies in the specimen, which in turn indicate a prior exposure of the patient to *Chlamydia*.

In another embodiment, antibodies that specifically recognize a *Chlamydia* peptide encoded by the nucleotide sequences disclosed herein are useful in diagnosing the presence of infection-specific *Chlamydia* antigens in a subject or sample. For example, detection of infection-specific antigens that are present in cells or tissues may be done by immunofluorescence, indirect-immunofluorescence and immunohistochemistry. In immunofluorescence, a fluorescent dye is bound directly to the antibody. In indirect-immunofluorescence, the dye is bound to an anti-immunoglobulin. Specific binding occurs between antigen and bound antibody is detected by virtue of fluorescent emissions from the dye moiety. This technique may be particularly useful, for instance, for detection of *Chlamydia* antigen present on a urogenital mucosal smear. *Chlamydia* may be present in urogenital mucosa, and a smear on a glass slide may be fixed and bathed in a solution containing an antibody specific to the infection-specific antigen. The slide is then washed to remove the unbound antibody, and a fluorescent anti-immunoglobulin antibody is added. The slide is washed again, and viewed microscopically under an appropriate wavelength of light to detect fluorescence. Fluorescence indicates the presence of *Chlamydia* antigen. Alternatively, a urogenital mucosal smear may be taken, the sample cultured with HeLa cells to produce large amounts of the RB form, and immunofluorescence may then be used to detect infection-specific *Chlamydia* antibodies.

Another aspect of the invention includes the use of nucleic acid primers to detect the presence of *Chlamydia* nucleic acids that encode infection-specific antigens in body samples and thus to diagnose infection. In other embodiments, these oligonucleotide primers will comprise at least 15 contiguous nucleotides of a DNA sequence as shown in SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, or 17. In other embodiments, such oligonucleotides may comprise at least 20 or at least 25 or more contiguous nucleotides of the aforementioned sequences.

One skilled in the art will appreciate that PCR primers are not required to exactly match the target gene sequence to which they anneal. Therefore, in another embodiment, the oligonucleotides will comprise a sequence of at least 15 nucleotides and preferably at least 20 nucleotides, the oligonucleotide sequence being substantially similar to a DNA sequence set forth

in SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, and 17. Such oligonucleotides may share at least about 75%, 85%, 90% or greater sequence identity.

The detection of specific nucleic acid sequences in a sample by polymerase chain reaction amplification (PCR) is discussed in detail in Innis et al., (1990). *PCR Protocols: A Guide to Methods and Applications*, Academic Press: San Diego, part 4 in particular. To detect *Chlamydia* sequences, primers based on the sequences disclosed herein would be synthesized, such that PCR amplification of a sample containing *Chlamydia* DNA would result in an amplified fragment of a predicted size. If necessary, the presence of this fragment following amplification of the sample nucleic acid could be detected by dot blot analysis. PCR amplification employing primers based on the sequences disclosed herein may also be employed to quantify the amounts of *Chlamydia* nucleic acid present in a particular sample (see chapters 8 and 9 of Innis et al., (1990)).

Alternatively, probes based on the nucleic acid sequences described herein may be labeled with suitable labels (such as P³² or biotin) and used in hybridization assays to detect the presence of *Chlamydia* nucleic acid in provided samples.

Reverse-transcription PCR using these primers may also be utilized to detect the presence of *Chlamydia* RNA which is indicative of an ongoing infection.

EXAMPLE 5: Production of Chlamydia Vaccines

The purified peptides of the present invention may be used directly as immunogens for vaccination. Methods for using purified peptides as vaccines are well known in the art and are described in Yang et al. (1991), Andersen (1994) and Jardim et al. (1990). As is well known in the art, adjuvants such as alum, Complete Freund's Adjuvant (CFA) and Incomplete Freund's Adjuvant (IFA) may be used in formulations of purified peptides as vaccines. Accordingly, one embodiment of the present invention is a vaccine comprising one or more immunostimulatory *C. trachomatis* or *C. psittaci* peptides encoded by nucleotide sequences as shown in the attached sequence listing, together with a pharmaceutically acceptable adjuvant.

Additionally a vaccine may comprise a defined fraction of the disclosed peptide of *C. trachomatis* or *C. psittaci* or may comprise a peptide wherein the gene coding for the peptide shows substantial similarity to the DNA sequences disclosed herein, such as for orthologous genes of *C. pneumoniae* or *C. pecorum*.

Additionally, the vaccines may be formulated using a peptide according to the present invention together with a pharmaceutically acceptable excipient such as water, saline, dextrose and glycerol. The vaccines may also include auxiliary substances such as emulsifying agents and pH buffers.

It will be appreciated by one of skill in the art that vaccines formulated as described above may be administered in a number of ways including subcutaneous, intra-muscular and intra-venous injection. Doses of the vaccine administered will vary depending on the antigenicity of the particular peptide or peptide combination employed in the vaccine, and characteristics of the

animal or human patient to be vaccinated. While the determination of individual doses will be within the skill of the administering physician, it is anticipated that doses of between 1 microgram and 1 milligram will be employed.

5 As with many vaccines, the vaccines of the present invention may routinely be administered several times over the course of a number of weeks to ensure that an effective immune response is triggered. Where such multiple doses are administered, they will normally be administered at from two to twelve week intervals, more usually from three to five week intervals. Periodic boosters at intervals of 1-5 years, usually three years, may be desirable to maintain the desired levels of protective immunity.

10 Alternatively, multiple immunostimulatory peptides may also be administered by expressing the nucleic acids encoding the peptides in a nonpathogenic microorganism, and using this transformed nonpathogenic microorganism as a vaccine.

15 Finally, a recent development in the field of vaccines is the direct injection of nucleic acid molecules encoding peptide antigens, as described in Janeway & Travers, (1997). Thus, plasmids which include nucleic acid molecules described herein, or which include nucleic acid sequences encoding peptides according to the present invention may be utilized in such DNA vaccination methods.

20 The vaccine of the invention may be used to inoculate potential animal targets of any of the chlamydial diseases including those caused by *C. trachomatis*, *C. psittaci*, *C. pneumoniae* or *C. pecorum*. Indeed the vaccine of the invention may be used to inoculate animals against any disease that shows immunological cross-protection as a result of exposure to infection-specific *Chlamydia* antigen. The protein or polypeptide is present in the vaccine in an amount sufficient to induce a protective immune response whether through humoral or cell mediated pathways or through both. Such a response protects the immunized animal against chlamydial infections
25 specifically by raising an immune response against the Reticulate Body form of *Chlamydia*.

The above embodiments are set out only by way of example and are not intended to be exclusive, one skilled in the art will understand that the invention may be practiced in various additional ways without departing from the subject of the spirit of the invention.

REFERENCES

- Akins, D. R., et al. (1997) *J. Bacteriol.* 179:5076-5086.
- Amann and Brosius (1985). *Gene* 40:183.
- Andersen (1994). *Infection & Immunity* 62:2536.
- 5 Ausubel et al. (1987). *Current Protocols in Molecular Biology*, ed. Greene Publishing and Wiley-Interscience: New York (with periodic updates).
- Blanco, D. R., et al. (1995) *J. Bacteriol.* 177:3556-3562.
- Bannantine, J.P., et al. (1997) *Abstr. Gen. Mtg. Amer. Soc. Microbiol.* D-004. Miami, FL.
- Blanco, D. R., et al. (1996) *J. Bacteriol.* 178:6685-6692.
- 10 Brown, W.J., and Farquhar, M.G. (1989) *Meth Cell Biol* 31:553-569.
- Caldwell, H.D., et al. (1981) *Infect. Immunol.* 31:1161-1176.
- Chou, P.Y. and Fasman, G.D. (1978) *Annu Rev Biochem* 47:251-276.
- Engvall (1980). *Enzymol.* 70:419.
- Gray et al. (1982). *Proc. Natl. Acad. Sci. USA* 79:6598.
- 15 Hackstadt, T., R. et al. (1992) *Infect. Immun.* 60:159-165.
- Hardham, J.M., et al. (1977) *Gene* 197:47-64.
- Harlow and Lane (1988). *Antibodies, A Laboratory Manual*, Cold Spring Harbor Laboratory, New York.
- Innis et al. (1990) *PCR Protocols: A Guide to Methods and Applications*, Academic Press: San
- 20 Diego.
- Janeway & Travers (1997) *Immunobiology: The Immune System in Health and Disease* 13.21. Garland Publishing, Inc. New York.
- Kohler and Milstein (1975) *Nature* 256:495.
- Kyte, J. and Doolittle, R.F. (1982) *J Mol Biol* 157:105-132.
- 25 Peeling, R. and Burnham, R. (1996) *Emerging Infectious Diseases* 2 (4) 307-317.
- Rockey, D.D., and Rosquist, J.L. (1994) *Infect Immun* 62:106-112.
- Rockey, D.D., et al. (1995) *Mol Microbiol* 15:617-626.
- Rockey, D.D., et al. (1996) *Infect Immun* 64:4269-4278.
- 30 Rockey, D.D., et al. (1997). *Mol Microbiol* 24:217-228.
- Robson, B. and Suzuki, E. (1976) *J Mol Biol* 107:327-356.
- Rockey, D.D., and Rosquist, J.L. (1994). *Infect Immun* 62:106-112.
- Rockey, D. D., et al. (1995) *Mol. Microbiol.* 15:617-626.
- Rockey, D.D., et al. (1997) *Mol. Microbiol.* 24:217-228.
- 35 Rothman, J.E., and F. T. Wieland (1996) *Science* 272:227-234.
- Ruther and Muller-Hill (1983). *EMBO J.* 2:1791.
- Sambrook et al. (1989) *Molecular Cloning: A Laboratory Manual*, 2nd ed., vol. 1-3, ed. Cold Spring Harbor Lab. Press: Cold Spring Harbor, NY.

- Shimatake and Rosenberg (1981). *Nature* (London) 292:128.
- Stanley and Luzio (1984). *EMBO J.* 3:1429.
- Studiar and Moffatt (1986). *J. Mol. Biol.* 189:113.
- Su, H., et al. (1990) *J. Exp. Med.* 172:203-212.
- 5 Yang et al. (1990) *J. Immunology* 145:2281-2285.
- Yuan, Y., et al. (1992) *Infect Immun* 60: 2288-2296.

CLAIMS

What is claimed is:

1. A purified infection-specific protein comprising an amino acid sequence selected from the group consisting of:
 - 5 (a) SEQ ID NO: 2,
 - (b) SEQ ID NO: 4,
 - (c) SEQ ID NO: 6,
 - (d) SEQ ID NO: 10,
 - (e) SEQ ID NO: 12,
 - 10 (f) an amino acid sequence that differs from an amino acid sequence of (a) to (e) inclusive, by one or more conservative amino acid substitutions, and
 - (g) an amino acid sequence having at least 60% sequence identity to an amino acid sequence of (a) to (e) inclusive.
2. An isolated nucleic acid molecule encoding a protein according to claim 1.
- 15 3. An isolated nucleic acid molecule according to claim 2 wherein the nucleic acid molecule comprises a nucleic acid sequence selected from the group consisting of:
 - (a) SEQ ID NO: 1,
 - (b) SEQ ID NO: 3,
 - (c) SEQ ID NO: 5,
 - 20 (d) SEQ ID NO: 9, and
 - (e) SEQ ID NO: 11.
4. A recombinant nucleic acid molecule comprising a promoter sequence operably linked to a nucleotide molecule according to claim 2.
5. A vaccine preparation comprising at least one purified peptide comprising at least
25 5 contiguous amino acids selected from the group consisting of:
 - (a) SEQ ID NO: 2,
 - (b) SEQ ID NO: 4,
 - (c) SEQ ID NO: 6,
 - (d) SEQ ID NO: 8,
 - 30 (e) SEQ ID NO: 10,
 - (f) SEQ ID NO: 12,
 - (g) SEQ ID NO: 14,
 - (h) SEQ ID NO: 16, and
 - (i) SEQ ID NO: 18.
- 35 6. The vaccine preparation of claim 5 wherein the peptide comprises at least 10 contiguous amino acids of at least one of the specified sequences.
7. The vaccine preparation of claim 5 wherein the peptide comprises at least 15 contiguous amino acids of at least one of the specified sequences.

8. The vaccine preparation of claim 5 wherein the purified peptide comprises at least 20 contiguous amino acids of at least one of the specified sequences.

9. A vaccine preparation comprising an amino acid sequence selected from the group consisting of:
- (a) SEQ ID NO: 2,
 - (b) SEQ ID NO: 4,
 - (c) SEQ ID NO: 6,
 - (d) SEQ ID NO: 8,
 - (e) SEQ ID NO: 10,
 - (f) SEQ ID NO: 12,
 - (g) SEQ ID NO: 14,
 - (h) SEQ ID NO: 16,
 - (i) SEQ ID NO: 18,
 - (j) an amino acid sequence that differs from an amino acid sequence of (a) to (i) inclusive, by one or more conservative amino acid substitutions, and
 - (k) an amino acid sequence having at least 60% sequence identity to an amino acid sequence of (a) to (i) inclusive.

10. A method of making a vaccine comprising combining a pharmaceutically acceptable excipient with a purified peptide having an amino acid sequence selected from the group consisting of:
- (a) SEQ ID NO:2,
 - (b) SEQ ID NO:4,
 - (c) SEQ ID NO:6,
 - (d) SEQ ID NO:8,
 - (e) SEQ ID NO:10,
 - (f) SEQ ID NO:12,
 - (g) SEQ ID NO:14,
 - (h) SEQ ID NO:16,
 - (i) SEQ ID NO:18,
 - (j) an amino acid sequence that differs from an amino acid sequence of (a) to (i) inclusive, by one or more conservative amino acid substitutions,
 - (k) an amino acid sequence having at least 60% sequence identity to an amino acid sequence of (a) to (i) inclusive, and
 - (l) at least 10 contiguous amino acids from an amino acid sequence of (a) to (i) inclusive.
11. A method of vaccination, comprising administering a vaccine preparation according to claim 5 to a mammal.

12. A method of vaccination, comprising administering a vaccine preparation according to claim 9 to a mammal.

13. A method of detecting an infection-specific *Chlamydia* protein in a biological sample comprising: contacting the biological sample with at least one anti-*Chlamydia* antibody, which antibody is an infection-specific antibody, such that a reaction between the antibody and the infection-specific *Chlamydia* protein gives rise to a detectable effect, and detecting the detectable effect.

14. The method of claim 13 wherein the anti-*Chlamydia* antibody binds specifically to a peptide having an amino acid sequence selected from the group consisting of:

- (a) SEQ ID NO: 2,
- (b) SEQ ID NO: 4,
- (c) SEQ ID NO: 6,
- (d) SEQ ID NO: 8,
- (e) SEQ ID NO: 10,
- (f) SEQ ID NO: 12,
- (g) SEQ ID NO: 14,
- (h) SEQ ID NO: 16, and
- (i) SEQ ID NO: 18.

15. A method of detecting an infection-specific anti-*Chlamydia* antibody in a biological sample comprising: contacting the biological sample with at least one *Chlamydia* peptide, which peptide is an infection specific peptide, such that a reaction between the peptide and the infection-specific anti-*Chlamydia* antibody gives rise to a detectable effect, and detecting the detectable effect.

16. The method of claim 15 wherein the *Chlamydia* peptide comprises at least 5 contiguous amino acids of a sequence selected from the group consisting of:

- (a) SEQ ID NO: 2,
- (b) SEQ ID NO: 4,
- (c) SEQ ID NO: 6,
- (d) SEQ ID NO: 8,
- (e) SEQ ID NO: 10,
- (f) SEQ ID NO: 12,
- (g) SEQ ID NO: 14,
- (h) SEQ ID NO: 16, and
- (i) SEQ ID NO: 18.

17. The method of claim 15 wherein said *Chlamydia* peptide comprises an amino acid sequence selected from the group consisting of:

- (a) SEQ ID NO: 2,
- (b) SEQ ID NO: 4,

-30-

- (c) SEQ ID NO: 6,
- (d) SEQ ID NO: 8,
- (e) SEQ ID NO: 10,
- (f) SEQ ID NO: 12,
- 5 (g) SEQ ID NO: 14,
- (h) SEQ ID NO: 16, and
- (i) SEQ ID NO: 18.

18. A method of treating a *Chlamydial* infection comprising directing a therapeutic agent against a specific target, said target chosen from the group consisting of: (i) an infection-specific protein of *Chlamydia*, (ii) a gene that encodes an infection-specific protein of *Chlamydia* and (iii) an RNA transcript that encodes an infection-specific protein of *Chlamydia*, wherein said therapeutic agent interacts with said target to affect a reduction in pathology.
- 10

SEQUENCE LISTING

<110> Oregon State University

<120> Methods of use for infection-specific INCA, INCB, and INCC proteins of Chlamydia

<130> 52297

<140>

<141>

<150> 60/082,588

<151> 1998-04-21

<150> 60/082,438

<151> 1998-04-20

<150> 60/086,450

<151> 1998-05-22

<160> 24

<170> PatentIn Ver. 2.0

<210> 1

<211> 534

<212> DNA

<213> Chlamydia trachomatis

<220>

<221> CDS

<222> (1)..(534)

<400> 1

atg	aaa	aag	ttc	tta	tta	ctt	agc	tta	atg	tct	ttg	tca	tct	cta	cct	48
Met	Lys	Lys	Phe	Leu	Leu	Leu	Ser	Leu	Met	Ser	Leu	Ser	Ser	Leu	Pro	
1				5					10					15		

aca	ttt	gca	gct	aat	tct	aca	ggc	aca	att	gga	atc	gtt	aat	tta	cgt	96
Thr	Phe	Ala	Ala	Asn	Ser	Thr	Gly	Thr	Ile	Gly	Ile	Val	Asn	Leu	Arg	
			20					25					30			

cgc	tgc	cta	gaa	gag	tct	gct	ctt	ggg	aaa	aaa	gaa	tct	gct	gaa	ttc	144
Arg	Cys	Leu	Glu	Glu	Ser	Ala	Leu	Gly	Lys	Lys	Glu	Ser	Ala	Glu	Phe	
		35					40					45				

gaa	aag	atg	aaa	aac	caa	ttc	tct	aac	agc	atg	ggg	aag	atg	gag	gaa	192
Glu	Lys	Met	Lys	Asn	Gln	Phe	Ser	Asn	Ser	Met	Gly	Lys	Met	Glu	Glu	
	50					55					60					

gaa	ctg	tct	tct	atc	tat	tcc	aag	ctc	caa	gac	gac	gat	tac	atg	gaa	240
Glu	Leu	Ser	Ser	Ile	Tyr	Ser	Lys	Leu	Gln	Asp	Asp	Asp	Tyr	Met	Glu	
	65				70					75					80	

ggc	cta	tcc	gag	acc	gca	gct	gcc	gaa	tta	aga	aaa	aaa	ttc	gaa	gat	288
Gly	Leu	Ser	Glu	Thr	Ala	Ala	Ala	Glu	Leu	Arg	Lys	Lys	Phe	Glu	Asp	
			85					90						95		

cta	tct	gca	gaa	tac	aac	aca	gct	caa	ggg	cag	tat	tac	caa	ata	tta	336
Leu	Ser	Ala	Glu	Tyr	Asn	Thr	Ala	Gln	Gly	Gln	Tyr	Tyr	Gln	Ile	Leu	

	100	105	110	
aac caa agt aat ttc aag cgc atg caa aag att atg gaa gaa gtg aaa				384
Asn Gln Ser Asn Phe Lys Arg Met Gln Lys Ile Met Glu Glu Val Lys				
	115	120	125	
aaa gct tct gaa act gtg cgt att caa gaa ggc ttg tca gtc ctt ctt				432
Lys Ala Ser Glu Thr Val Arg Ile Gln Glu Gly Leu Ser Val Leu Leu				
	130	135	140	
aac gaa gat att gtc tta tct atc gat agt tcg gca gat aaa acc gat				480
Asn Glu Asp Ile Val Leu Ser Ile Asp Ser Ser Ala Asp Lys Thr Asp				
	145	150	155	160
gct gtt att aaa gtt ctt gat gtt ctt ttc aaa ata att aac atg cga				528
Ala Val Ile Lys Val Leu Asp Val Leu Phe Lys Ile Ile Asn Met Arg				
	165	170	175	
agc tag				534
Ser				

<210> 2

<211> 177

<212> PRT

<213> Chlamydia trachomatis

<400> 2

Met Lys Lys Phe Leu Leu Leu Ser Leu Met Ser Leu Ser Ser Leu Pro				
1	5	10	15	
Thr Phe Ala Ala Asn Ser Thr Gly Thr Ile Gly Ile Val Asn Leu Arg				
20	25	30		
Arg Cys Leu Glu Glu Ser Ala Leu Gly Lys Lys Glu Ser Ala Glu Phe				
35	40	45		
Glu Lys Met Lys Asn Gln Phe Ser Asn Ser Met Gly Lys Met Glu Glu				
50	55	60		
Glu Leu Ser Ser Ile Tyr Ser Lys Leu Gln Asp Asp Asp Tyr Met Glu				
65	70	75	80	
Gly Leu Ser Glu Thr Ala Ala Ala Glu Leu Arg Lys Lys Phe Glu Asp				
85	90	95		
Leu Ser Ala Glu Tyr Asn Thr Ala Gln Gly Gln Tyr Tyr Gln Ile Leu				
100	105	110		
Asn Gln Ser Asn Phe Lys Arg Met Gln Lys Ile Met Glu Glu Val Lys				
115	120	125		
Lys Ala Ser Glu Thr Val Arg Ile Gln Glu Gly Leu Ser Val Leu Leu				
130	135	140		
Asn Glu Asp Ile Val Leu Ser Ile Asp Ser Ser Ala Asp Lys Thr Asp				
145	150	155	160	
Ala Val Ile Lys Val Leu Asp Val Leu Phe Lys Ile Ile Asn Met Arg				
165	170	175		

Ser

<210> 3

<211> 846

<212> DNA

<213> Chlamydia trachomatis

<220>

<221> CDS

<222> (1)..(846)

<400> 3

atg aat cgc atg att tgt gat tgc gtg tct cgc ata act ggg gat cga	48
Met Asn Arg Met Ile Cys Asp Cys Val Ser Arg Ile Thr Gly Asp Arg	
1 5 10 15	
gtc aag aat att gtt ctg att gat gga gcg att gat cct cat tca tat	96
Val Lys Asn Ile Val Leu Ile Asp Gly Ala Ile Asp Pro His Ser Tyr	
20 25 30	
gag atg gtg aag ggg gat gaa gac cga atg gct atg agc cag ctg att	144
Glu Met Val Lys Gly Asp Glu Asp Arg Met Ala Met Ser Gln Leu Ile	
35 40 45	
ttt tgc aat ggt tta ggt tta gag cat tca gct agt tta cgt aaa cat	192
Phe Cys Asn Gly Leu Gly Leu Glu His Ser Ala Ser Leu Arg Lys His	
50 55 60	
cta gag ggt aac cca aaa gtc gtt gat tta ggt caa cgt ttg ctt aac	240
Leu Glu Gly Asn Pro Lys Val Val Asp Leu Gly Gln Arg Leu Leu Asn	
65 70 75 80	
aaa aac tgt ttt gat ctt ctg agt gaa gaa gga ttc cct gac cca cat	288
Lys Asn Cys Phe Asp Leu Leu Ser Glu Glu Gly Phe Pro Asp Pro His	
85 90 95	
att tgg acg gat atg aga gta tgg ggt gct gct gta aaa gag atg gct	336
Ile Trp Thr Asp Met Arg Val Trp Gly Ala Ala Val Lys Glu Met Ala	
100 105 110	
gcg gca tta att caa caa ttt cct caa tat gaa gaa gat ttt caa aag	384
Ala Ala Leu Ile Gln Gln Phe Pro Gln Tyr Glu Glu Asp Phe Gln Lys	
115 120 125	
aat gcg gat cag atc tta tca gag atg gag gaa ctt gat cgt tgg gca	432
Asn Ala Asp Gln Ile Leu Ser Glu Met Glu Glu Leu Asp Arg Trp Ala	
130 135 140	
gtg cgt tct ctc tct acg att cct gaa aaa aat cgc tat tta gtc aca	480
Val Arg Ser Leu Ser Thr Ile Pro Glu Lys Asn Arg Tyr Leu Val Thr	
145 150 155 160	
ggc cac aat gcg ttc agt tac ttt act cgt cgg tat cta tcc tct gat	528
Gly His Asn Ala Phe Ser Tyr Phe Thr Arg Arg Tyr Leu Ser Ser Asp	
165 170 175	
gcg gag aga gtg tct ggg gaa tgg aga tcg cgt tgc att tct cca gaa	576
Ala Glu Arg Val Ser Gly Glu Trp Arg Ser Arg Cys Ile Ser Pro Glu	
180 185 190	

```

ggg ttg tct cct gag gct cag att agt atc cga gat att atg cgt gta 624
Gly Leu Ser Pro Glu Ala Gln Ile Ser Ile Arg Asp Ile Met Arg Val
      195                      200                      205

gtg gag tat atc tct gca aac gat gta gaa gtt gtc ttt tta gag gat 672
Val Glu Tyr Ile Ser Ala Asn Asp Val Glu Val Val Phe Leu Glu Asp
      210                      215                      220

acg tta aat caa gat gct ttg aga aag att gtt tct tgc tct aag agc 720
Thr Leu Asn Gln Asp Ala Leu Arg Lys Ile Val Ser Cys Ser Lys Ser
      225                      230                      235                      240

gga caa aag att cgt ctc gct aag tct cct tta tat agc gat aat gtc 768
Gly Gln Lys Ile Arg Leu Ala Lys Ser Pro Leu Tyr Ser Asp Asn Val
      245                      250                      255

tgt gat aac tat ttt agc acg ttc cag cac aat gtt cgc aca att aca 816
Cys Asp Asn Tyr Phe Ser Thr Phe Gln His Asn Val Arg Thr Ile Thr
      260                      265                      270

gaa gaa ttg gga ggg act gtt ctt gaa tag 846
Glu Glu Leu Gly Gly Thr Val Leu Glu
      275                      280

```

<210> 4

<211> 281

<212> PRT

<213> Chlamydia trachomatis

<400> 4

```

Met Asn Arg Met Ile Cys Asp Cys Val Ser Arg Ile Thr Gly Asp Arg
  1                      5                      10                      15

Val Lys Asn Ile Val Leu Ile Asp Gly Ala Ile Asp Pro His Ser Tyr
      20                      25                      30

Glu Met Val Lys Gly Asp Glu Asp Arg Met Ala Met Ser Gln Leu Ile
      35                      40                      45

Phe Cys Asn Gly Leu Gly Leu Glu His Ser Ala Ser Leu Arg Lys His
      50                      55                      60

Leu Glu Gly Asn Pro Lys Val Val Asp Leu Gly Gln Arg Leu Leu Asn
      65                      70                      75                      80

Lys Asn Cys Phe Asp Leu Leu Ser Glu Glu Gly Phe Pro Asp Pro His
      85                      90                      95

Ile Trp Thr Asp Met Arg Val Trp Gly Ala Ala Val Lys Glu Met Ala
      100                      105                      110

Ala Ala Leu Ile Gln Gln Phe Pro Gln Tyr Glu Glu Asp Phe Gln Lys
      115                      120                      125

Asn Ala Asp Gln Ile Leu Ser Glu Met Glu Glu Leu Asp Arg Trp Ala
      130                      135                      140

Val Arg Ser Leu Ser Thr Ile Pro Glu Lys Asn Arg Tyr Leu Val Thr
      145                      150                      155                      160

```

Gly His Asn Ala Phe Ser Tyr Phe Thr Arg Arg Tyr Leu Ser Ser Asp
 165 170 175

Ala Glu Arg Val Ser Gly Glu Trp Arg Ser Arg Cys Ile Ser Pro Glu
 180 185 190

Gly Leu Ser Pro Glu Ala Gln Ile Ser Ile Arg Asp Ile Met Arg Val
 195 200 205

Val Glu Tyr Ile Ser Ala Asn Asp Val Glu Val Val Phe Leu Glu Asp
 210 215 220

Thr Leu Asn Gln Asp Ala Leu Arg Lys Ile Val Ser Cys Ser Lys Ser
 225 230 235 240

Gly Gln Lys Ile Arg Leu Ala Lys Ser Pro Leu Tyr Ser Asp Asn Val
 245 250 255

Cys Asp Asn Tyr Phe Ser Thr Phe Gln His Asn Val Arg Thr Ile Thr
 260 265 270

Glu Glu Leu Gly Gly Thr Val Leu Glu
 275 280

<210> 5

<211> 861

<212> DNA

<213> Chlamydia trachomatis

<220>

<221> CDS

<222> (1)..(861)

<400> 5

atg tct gtg ata act att tta gca cgt tcc agc aca atg ttc gca caa	48
Met Ser Val Ile Thr Ile Leu Ala Arg Ser Ser Thr Met Phe Ala Gln	
1 5 10 15	
tta cag aag aat tgg gag gga ctg ttc ttg aat aga gat aat gca att	96
Leu Gln Lys Asn Trp Glu Gly Leu Phe Leu Asn Arg Asp Asn Ala Ile	
20 25 30	
gct tgg tcc gta gag gat ctt tgt gtt aat tat gat cac tca gac gtc	144
Ala Trp Ser Val Glu Asp Leu Cys Val Asn Tyr Asp His Ser Asp Val	
35 40 45	
tta tgt cac att act ttt tct ctg cct gca ggg gca atg gct gct att	192
Leu Cys His Ile Thr Phe Ser Leu Pro Ala Gly Ala Met Ala Ala Ile	
50 55 60	
att ggg ccg aat gga gct ggt aaa agt act ttg ctt aag gct tct tta	240
Ile Gly Pro Asn Gly Ala Gly Lys Ser Thr Leu Leu Lys Ala Ser Leu	
65 70 75 80	
gga ctg att cgt gct tct tct ggc caa agc ttg ttc ttt ggt cag aga	288
Gly Leu Ile Arg Ala Ser Ser Gly Gln Ser Leu Phe Phe Gly Gln Arg	
85 90 95	
ttt tcc aag gca cat cat aga ata gcc tat atg cct caa aga gcg agt	336
Phe Ser Lys Ala His His Arg Ile Ala Tyr Met Pro Gln Arg Ala Ser	

100	105	110	
gtg gat tgg gat ttc cca atg act gtt ctt gat ctc gtg ttg atg ggg			384
Val Asp Trp Asp Phe Pro Met Thr Val Leu Asp Leu Val Leu Met Gly			
115	120	125	
tgt tac ggc tat aaa gga ata tgg aat cgt att tcc act gat gat cgt			432
Cys Tyr Gly Tyr Lys Gly Ile Trp Asn Arg Ile Ser Thr Asp Asp Arg			
130	135	140	
cag gag gct atg cgt att tta gag cgg gtt ggt ttg gaa gct ttt gca			480
Gln Glu Ala Met Arg Ile Leu Glu Arg Val Gly Leu Glu Ala Phe Ala			
145	150	155	160
aat cgt caa ata ggt aag ctc tct gga gga caa caa cag aga gct ttt			528
Asn Arg Gln Ile Gly Lys Leu Ser Gly Gly Gln Gln Gln Arg Ala Phe			
165	170	175	
tta gcg cgg tca tta atg caa aaa gca gat ttg tat ctc atg gat gag			576
Leu Ala Arg Ser Leu Met Gln Lys Ala Asp Leu Tyr Leu Met Asp Glu			
180	185	190	
ctg ttc tct gcg atc gat atg gcc tct tat cag atg gtt gta gat gtt			624
Leu Phe Ser Ala Ile Asp Met Ala Ser Tyr Gln Met Val Val Asp Val			
195	200	205	
ttg caa gag ctt aaa agc gaa ggg aag act att gtg gtc att cat cat			672
Leu Gln Glu Leu Lys Ser Glu Gly Lys Thr Ile Val Val Ile His His			
210	215	220	
gat ttg agt aat gtc cgg aag ctt ttt gat cat gtg att tta tta aat			720
Asp Leu Ser Asn Val Arg Lys Leu Phe Asp His Val Ile Leu Leu Asn			
225	230	235	240
aag cat ctt gtg tgc tct gga agc gta gaa gaa tgc ttg act aaa gaa			768
Lys His Leu Val Cys Ser Gly Ser Val Glu Glu Cys Leu Thr Lys Glu			
245	250	255	
gcc att ttt cag gct tat ggg tgt gac ttg agc ttt tgg att aca cac			816
Ala Ile Phe Gln Ala Tyr Gly Cys Asp Leu Ser Phe Trp Ile Thr His			
260	265	270	
tca aat tgt cta gag gca agt acc aag gat cgt gct aga tgc tga			861
Ser Asn Cys Leu Glu Ala Ser Thr Lys Asp Arg Ala Arg Cys			
275	280	285	

<210> 6

<211> 286

<212> PRT

<213> Chlamydia trachomatis

<400> 6

Met	Ser	Val	Ile	Thr	Ile	Leu	Ala	Arg	Ser	Ser	Thr	Met	Phe	Ala	Gln
1					5				10					15	

Leu	Gln	Lys	Asn	Trp	Glu	Gly	Leu	Phe	Leu	Asn	Arg	Asp	Asn	Ala	Ile
		20						25					30		

Ala	Trp	Ser	Val	Glu	Asp	Leu	Cys	Val	Asn	Tyr	Asp	His	Ser	Asp	Val
		35					40					45			

Leu Cys His Ile Thr Phe Ser Leu Pro Ala Gly Ala Met Ala Ala Ile
 50 55 60
 Ile Gly Pro Asn Gly Ala Gly Lys Ser Thr Leu Leu Lys Ala Ser Leu
 65 70 75 80
 Gly Leu Ile Arg Ala Ser Ser Gly Gln Ser Leu Phe Phe Gly Gln Arg
 85 90 95
 Phe Ser Lys Ala His His Arg Ile Ala Tyr Met Pro Gln Arg Ala Ser
 100 105 110
 Val Asp Trp Asp Phe Pro Met Thr Val Leu Asp Leu Val Leu Met Gly
 115 120 125
 Cys Tyr Gly Tyr Lys Gly Ile Trp Asn Arg Ile Ser Thr Asp Asp Arg
 130 135 140
 Gln Glu Ala Met Arg Ile Leu Glu Arg Val Gly Leu Glu Ala Phe Ala
 145 150 155 160
 Asn Arg Gln Ile Gly Lys Leu Ser Gly Gly Gln Gln Gln Arg Ala Phe
 165 170 175
 Leu Ala Arg Ser Leu Met Gln Lys Ala Asp Leu Tyr Leu Met Asp Glu
 180 185 190
 Leu Phe Ser Ala Ile Asp Met Ala Ser Tyr Gln Met Val Val Asp Val
 195 200 205
 Leu Gln Glu Leu Lys Ser Glu Gly Lys Thr Ile Val Val Ile His His
 210 215 220
 Asp Leu Ser Asn Val Arg Lys Leu Phe Asp His Val Ile Leu Leu Asn
 225 230 235 240
 Lys His Leu Val Cys Ser Gly Ser Val Glu Glu Cys Leu Thr Lys Glu
 245 250 255
 Ala Ile Phe Gln Ala Tyr Gly Cys Asp Leu Ser Phe Trp Ile Thr His
 260 265 270
 Ser Asn Cys Leu Glu Ala Ser Thr Lys Asp Arg Ala Arg Cys
 275 280 285

<210> 7

<211> 1068

<212> DNA

<213> Chlamydia psittaci

<220>

<221> CDS

<222> (1)..(1068)

<400> 7

atg aca gta tcc aca gac aac aca agt cct gta ata tcg aga gcg tcc 48
 Met Thr Val Ser Thr Asp Asn Thr Ser Pro Val Ile Ser Arg Ala Ser
 1 5 10 15

tca cct act ttt gga gat cat ggt aag gat ttc gac aac aat aaa att	96
Ser Pro Thr Phe Gly Asp His Gly Lys Asp Phe Asp Asn Asn Lys Ile	
20 25 30	
ata ccc att tca ata gaa gct cca act tct tca gct gct gct gta ggg	144
Ile Pro Ile Ser Ile Glu Ala Pro Thr Ser Ser Ala Ala Ala Val Gly	
35 40 45	
gct aaa acg gct atc gag cct gaa gga aga agc cca cta ctt caa agg	192
Ala Lys Thr Ala Ile Glu Pro Glu Gly Arg Ser Pro Leu Leu Gln Arg	
50 55 60	
att tgc tat ctt gtt aaa att atc gct gcc atc gcc ctc ttt gtt gtt	240
Ile Cys Tyr Leu Val Lys Ile Ile Ala Ala Ile Ala Leu Phe Val Val	
65 70 75 80	
ggt atc gca gcc tta gtt tgc tta tat ctc ggt agc gtt atc tca acg	288
Gly Ile Ala Ala Leu Val Cys Leu Tyr Leu Gly Ser Val Ile Ser Thr	
85 90 95	
cct tct ctt att ctt atg ctt gcg atc atg ctt gta tcc ttt gtg atc	336
Pro Ser Leu Ile Leu Met Leu Ala Ile Met Leu Val Ser Phe Val Ile	
100 105 110	
gtt att acg gca att cga gat ggc aca ccg tct caa gtg gtc cgt cac	384
Val Ile Thr Ala Ile Arg Asp Gly Thr Pro Ser Gln Val Val Arg His	
115 120 125	
atg aaa cag caa att cag caa ttt ggc gaa gaa aac acg cgt tta cat	432
Met Lys Gln Gln Ile Gln Gln Phe Gly Glu Glu Asn Thr Arg Leu His	
130 135 140	
acc gca gta gaa aat cta aaa gct gtt aac gtt gag ctc tca gag caa	480
Thr Ala Val Glu Asn Leu Lys Ala Val Asn Val Glu Leu Ser Glu Gln	
145 150 155 160	
att aac caa ctt aaa caa cta cat act aga tta tcg gat ttt ggt gat	528
Ile Asn Gln Leu Lys Gln Leu His Thr Arg Leu Ser Asp Phe Gly Asp	
165 170 175	
agg ctt gaa gcg aat acc ggt gat ttt act gca ctt att gcg gat ttc	576
Arg Leu Glu Ala Asn Thr Gly Asp Phe Thr Ala Leu Ile Ala Asp Phe	
180 185 190	
caa ctc agt ctg gaa gag ttt aag tct gtt ggt act aaa gtt gaa acc	624
Gln Leu Ser Leu Glu Glu Phe Lys Ser Val Gly Thr Lys Val Glu Thr	
195 200 205	
atg ctc tct cca ttt gag aaa tta gct cag tct ttg aaa gag acc ttt	672
Met Leu Ser Pro Phe Glu Lys Leu Ala Gln Ser Leu Lys Glu Thr Phe	
210 215 220	
tct caa gaa gct gtt cag gca atg atg tcc tct gta act gag tta aga	720
Ser Gln Glu Ala Val Gln Ala Met Met Ser Ser Val Thr Glu Leu Arg	
225 230 235 240	
acc aat ttg aat gca ttg aaa gag ctt ata aca gag aat aaa acc gta	768
Thr Asn Leu Asn Ala Leu Lys Glu Leu Ile Thr Glu Asn Lys Thr Val	
245 250 255	
ata gag caa cta aaa gct gat gct caa ctt aga gaa gag caa gtg cgg	816

```

Ile Glu Gln Leu Lys Ala Asp Ala Gln Leu Arg Glu Glu Gln Val Arg
      260                      265                      270

ttt tta gaa aag cgt aaa caa gag tta gaa gag gct tgt tca aca ttg   864
Phe Leu Glu Lys Arg Lys Gln Glu Leu Glu Ala Cys Ser Thr Leu
      275                      280                      285

tcc cat tca att gcg act cta cag gaa tcc aca acc ctt cta aag gac   912
Ser His Ser Ile Ala Thr Leu Gln Glu Ser Thr Thr Leu Leu Lys Asp
      290                      295                      300

tct aca act aac tta cat gca gtt gaa agt cgt ctt atc ggt gtt atg   960
Ser Thr Thr Asn Leu His Ala Val Glu Ser Arg Leu Ile Gly Val Met
      305                      310                      315

gtt cag gat ggt gca gag tcc tcc acc gta gag gaa gct tca caa gat   1008
Val Gln Asp Gly Ala Glu Ser Ser Thr Val Glu Glu Ala Ser Gln Asp
      325                      330                      335

gat agc gcg caa ccc caa gat gaa aat caa tct gat gct gga gag cat   1056
Asp Ser Ala Gln Pro Gln Asp Glu Asn Gln Ser Asp Ala Gly Glu His
      340                      345                      350

aaa gat agt taa
Lys Asp Ser
      355

<210> 8
<211> 355
<212> PRT
<213> Chlamydia psittaci

<400> 8
Met Thr Val Ser Thr Asp Asn Thr Ser Pro Val Ile Ser Arg Ala Ser
  1                      5                      10                      15

Ser Pro Thr Phe Gly Asp His Gly Lys Asp Phe Asp Asn Asn Lys Ile
      20                      25                      30

Ile Pro Ile Ser Ile Glu Ala Pro Thr Ser Ser Ala Ala Ala Val Gly
      35                      40                      45

Ala Lys Thr Ala Ile Glu Pro Glu Gly Arg Ser Pro Leu Leu Gln Arg
      50                      55                      60

Ile Cys Tyr Leu Val Lys Ile Ile Ala Ala Ile Ala Leu Phe Val Val
      65                      70                      75                      80

Gly Ile Ala Ala Leu Val Cys Leu Tyr Leu Gly Ser Val Ile Ser Thr
      85                      90                      95

Pro Ser Leu Ile Leu Met Leu Ala Ile Met Leu Val Ser Phe Val Ile
      100                      105                      110

Val Ile Thr Ala Ile Arg Asp Gly Thr Pro Ser Gln Val Val Arg His
      115                      120                      125

Met Lys Gln Gln Ile Gln Gln Phe Gly Glu Glu Asn Thr Arg Leu His
      130                      135                      140

```

Thr Ala Val Glu Asn Leu Lys Ala Val Asn Val Glu Leu Ser Glu Gln
 145 150 155 160
 Ile Asn Gln Leu Lys Gln Leu His Thr Arg Leu Ser Asp Phe Gly Asp
 165 170 175
 Arg Leu Glu Ala Asn Thr Gly Asp Phe Thr Ala Leu Ile Ala Asp Phe
 180 185 190
 Gln Leu Ser Leu Glu Glu Phe Lys Ser Val Gly Thr Lys Val Glu Thr
 195 200 205
 Met Leu Ser Pro Phe Glu Lys Leu Ala Gln Ser Leu Lys Glu Thr Phe
 210 215 220
 Ser Gln Glu Ala Val Gln Ala Met Met Ser Ser Val Thr Glu Leu Arg
 225 230 235 240
 Thr Asn Leu Asn Ala Leu Lys Glu Leu Ile Thr Glu Asn Lys Thr Val
 245 250 255
 Ile Glu Gln Leu Lys Ala Asp Ala Gln Leu Arg Glu Glu Gln Val Arg
 260 265 270
 Phe Leu Glu Lys Arg Lys Gln Glu Leu Glu Glu Ala Cys Ser Thr Leu
 275 280 285
 Ser His Ser Ile Ala Thr Leu Gln Glu Ser Thr Thr Leu Leu Lys Asp
 290 295 300
 Ser Thr Thr Asn Leu His Ala Val Glu Ser Arg Leu Ile Gly Val Met
 305 310 315 320
 Val Gln Asp Gly Ala Glu Ser Ser Thr Val Glu Glu Ala Ser Gln Asp
 325 330 335
 Asp Ser Ala Gln Pro Gln Asp Glu Asn Gln Ser Asp Ala Gly Glu His
 340 345 350
 Lys Asp Ser
 355

<210> 9
 <211> 597
 <212> DNA
 <213> Chlamydia psittaci

<220>
 <221> CDS
 <222> (1)..(597)

<400> 9
 atg tca aca aca cca gca tct tca gca agt cga gac gta tta tta gat 48
 Met Ser Thr Thr Pro Ala Ser Ser Ala Ser Arg Asp Val Leu Leu Asp
 1 5 10 15
 gac gtt tta ata gct ttt aat aga aag cta aat ctc gta gaa caa caa 96
 Asp Val Leu Ile Ala Phe Asn Arg Lys Leu Asn Leu Val Glu Gln Gln
 20 25 30

gcg aaa gaa ctt gaa acg aaa gtc agt ttg gta gac aga aca gct act 144
 Ala Lys Glu Leu Glu Thr Lys Val Ser Leu Val Asp Arg Thr Ala Thr
 35 40 45

tta tca ctt acc act ggc aat aat gta gcc acg gat gta ctc ctt tta 192
 Leu Ser Leu Thr Thr Gly Asn Asn Val Ala Thr Asp Val Leu Leu Leu
 50 55 60

aaa gat gag gtt gca gaa cta aaa gga tgt ttg tct gca gtt acg gat 240
 Lys Asp Glu Val Ala Glu Leu Lys Gly Cys Leu Ser Ala Val Thr Asp
 65 70 75 80

cta tta atc cgc tca ggc tca tca aga aca cct ggg ggt gct cct aat 288
 Leu Leu Ile Arg Ser Gly Ser Ser Arg Thr Pro Gly Gly Ala Pro Asn
 85 90 95

cca gaa ggc act aat tac cta ata gga tgc aca cct cct tct ctt tgc 336
 Pro Glu Gly Thr Asn Tyr Leu Ile Gly Cys Thr Pro Pro Ser Leu Cys
 100 105 110

gct aaa ctt aca gcg tta gcg tta aca att ata gcc ctc att gct atc 384
 Ala Lys Leu Thr Ala Leu Ala Leu Thr Ile Ile Ala Leu Ile Ala Ile
 115 120 125

aca gta ctt gtt atc tgt att gtt act gtt tgc gcc ggt ttc ccc cta 432
 Thr Val Leu Val Ile Cys Ile Val Thr Val Cys Gly Gly Phe Pro Leu
 130 135 140

ttt att tcc cta ctc aac atg tac aca gtt ggt gct tgt ata tcc tta 480
 Phe Ile Ser Leu Leu Asn Met Tyr Thr Val Gly Ala Cys Ile Ser Leu
 145 150 155 160

ccg atc att tcg tgt gcc gca gtt tca atg atg att cta tgc tca cat 528
 Pro Ile Ile Ser Cys Ala Ala Val Ser Met Met Ile Leu Cys Ser His
 165 170 175

tct att aac tct tta tta aga aac agg cct gcg atc tat atg act aac 576
 Ser Ile Asn Ser Leu Leu Arg Asn Arg Pro Ala Ile Tyr Met Thr Asn
 180 185 190

aat ttt caa aca gaa tct taa 597
 Asn Phe Gln Thr Glu Ser
 195

<210> 10

<211> 198

<212> PRT

<213> Chlamydia psittaci

<400> 10

Met Ser Thr Thr Pro Ala Ser Ser Ala Ser Arg Asp Val Leu Leu Asp
 1 5 10 15

Asp Val Leu Ile Ala Phe Asn Arg Lys Leu Asn Leu Val Glu Gln Gln
 20 25 30

Ala Lys Glu Leu Glu Thr Lys Val Ser Leu Val Asp Arg Thr Ala Thr
 35 40 45

Leu Ser Leu Thr Thr Gly Asn Asn Val Ala Thr Asp Val Leu Leu Leu

50 55 60
 Lys Asp Glu Val Ala Glu Leu Lys Gly Cys Leu Ser Ala Val Thr Asp
 65 70 75 80
 Leu Leu Ile Arg Ser Gly Ser Ser Arg Thr Pro Gly Gly Ala Pro Asn
 85 90 95
 Pro Glu Gly Thr Asn Tyr Leu Ile Gly Cys Thr Pro Pro Ser Leu Cys
 100 105 110
 Ala Lys Leu Thr Ala Leu Ala Leu Thr Ile Ile Ala Leu Ile Ala Ile
 115 120 125
 Thr Val Leu Val Ile Cys Ile Val Thr Val Cys Gly Gly Phe Pro Leu
 130 135 140
 Phe Ile Ser Leu Leu Asn Met Tyr Thr Val Gly Ala Cys Ile Ser Leu
 145 150 155 160
 Pro Ile Ile Ser Cys Ala Ala Val Ser Met Met Ile Leu Cys Ser His
 165 170 175
 Ser Ile Asn Ser Leu Leu Arg Asn Arg Pro Ala Ile Tyr Met Thr Asn
 180 185 190
 Asn Phe Gln Thr Glu Ser
 195

<210> 11
 <211> 561
 <212> DNA
 <213> Chlamydia psittaci

<220>
 <221> CDS
 <222> (1)..(561)

<400> 11
 atg acc tct gta aga acc gat tta act cca ggc gac acc tca ctc caa 48
 Met Thr Ser Val Arg Thr Asp Leu Thr Pro Gly Asp Thr Ser Leu Gln
 1 5 10 15
 tct tct tta tta aat ccg agt gat ctc aca aca caa cta tcc aac ctc 96
 Ser Ser Leu Leu Asn Pro Ser Asp Leu Thr Thr Gln Leu Ser Asn Leu
 20 25 30
 cag act gtt ctc gca ggg ata caa caa caa cat cct tta aac ggt ggt 144
 Gln Thr Val Leu Ala Gly Ile Gln Gln Gln His Pro Leu Asn Gly Gly
 35 40 45
 tgg cct cag cat cat cct act ggc gct gca gat caa aat tat ctc atg 192
 Trp Pro Gln His His Pro Thr Gly Ala Ala Asp Gln Asn Tyr Leu Met
 50 55 60
 cgt ctg atg caa tct cat atg gca agt acc gta tca gca gta tct gaa 240
 Arg Leu Met Gln Ser His Met Ala Ser Thr Val Ser Ala Val Ser Glu
 65 70 75 80
 tta aga acc gaa gtc act gca atc aag aca aaa ttg cac ggg cta tct 288

Leu Arg Thr Glu Val Thr Ala Ile Lys Thr Lys Leu His Gly Leu Ser
 85 90 95

 act cca gct aat gtt tgc agc ggt cct atg gct cta gcc gct ttt ctt 336
 Thr Pro Ala Asn Val Cys Ser Gly Pro Met Ala Leu Ala Ala Phe Leu
 100 105 110

 cta gct ata tct tta gtt gcg att atc atc att gtt tta gcc tcc tta 384
 Leu Ala Ile Ser Leu Val Ala Ile Ile Ile Ile Val Leu Ala Ser Leu
 115 120 125

 ggc ctt gca ggc ata cta cct caa gct gcc gct atc tta gtg aat aca 432
 Gly Leu Ala Gly Ile Leu Pro Gln Ala Ala Ala Ile Leu Val Asn Thr
 130 135 140

 gca aac tct ata tgg gct att gtt agc gct tgc ata gtc act gtt atc 480
 Ala Asn Ser Ile Trp Ala Ile Val Ser Ala Ser Ile Val Thr Val Ile
 145 150 155 160

 tgc tta att agc gtg cta tgc ata acg cta att cga cac cat aaa ccc 528
 Cys Leu Ile Ser Val Leu Cys Ile Thr Leu Ile Arg His His Lys Pro
 165 170 175

 tta cct att gaa act agg cct acc gga cat taa 561
 Leu Pro Ile Glu Thr Arg Pro Thr Gly His
 180 185

 <210> 12
 <211> 186
 <212> PRT
 <213> Chlamydia psittaci

 <400> 12
 Met Thr Ser Val Arg Thr Asp Leu Thr Pro Gly Asp Thr Ser Leu Gln
 1 5 10 15

 Ser Ser Leu Leu Asn Pro Ser Asp Leu Thr Thr Gln Leu Ser Asn Leu
 20 25 30

 Gln Thr Val Leu Ala Gly Ile Gln Gln Gln His Pro Leu Asn Gly Gly
 35 40 45

 Trp Pro Gln His His Pro Thr Gly Ala Ala Asp Gln Asn Tyr Leu Met
 50 55 60

 Arg Leu Met Gln Ser His Met Ala Ser Thr Val Ser Ala Val Ser Glu
 65 70 75 80

 Leu Arg Thr Glu Val Thr Ala Ile Lys Thr Lys Leu His Gly Leu Ser
 85 90 95

 Thr Pro Ala Asn Val Cys Ser Gly Pro Met Ala Leu Ala Ala Phe Leu
 100 105 110

 Leu Ala Ile Ser Leu Val Ala Ile Ile Ile Ile Val Leu Ala Ser Leu
 115 120 125

 Gly Leu Ala Gly Ile Leu Pro Gln Ala Ala Ala Ile Leu Val Asn Thr
 130 135 140

Ala Asn Ser Ile Trp Ala Ile Val Ser Ala Ser Ile Val Thr Val Ile
 145 150 155 160

Cys Leu Ile Ser Val Leu Cys Ile Thr Leu Ile Arg His His Lys Pro
 165 170 175

Leu Pro Ile Glu Thr Arg Pro Thr Gly His
 180 185

<210> 13

<211> 822

<212> DNA

<213> Chlamydia trachomatis

<220>

<221> CDS

<222> (1)..(822)

<400> 13

atg aca acg cct act cta atc gtg att cct cca tct ccc cct gca cct 48
 Met Thr Thr Pro Thr Leu Ile Val Ile Pro Pro Ser Pro Pro Ala Pro
 1 5 10 15

tcc tac tca gcc aat cgc gta cct caa cct tct ttg atg gac aaa att 96
 Ser Tyr Ser Ala Asn Arg Val Pro Gln Pro Ser Leu Met Asp Lys Ile
 20 25 30

aag aaa ata gca gcc att gcc tcc cta att ctt ata ggc aca ata ggc 144
 Lys Lys Ile Ala Ala Ile Ala Ser Leu Ile Leu Ile Gly Thr Ile Gly
 35 40 45

ttt tta gct ctt ttg gga cat ctt gtt ggc ttt ctg atc gct cca caa 192
 Phe Leu Ala Leu Leu Gly His Leu Val Gly Phe Leu Ile Ala Pro Gln
 50 55 60

atc act att gtt ctt ctt gcc cta ttc att acc tca tta gca ggg aat 240
 Ile Thr Ile Val Leu Leu Ala Leu Phe Ile Thr Ser Leu Ala Gly Asn
 65 70 75 80

gct ctt tat cta cag aaa acc gct aat cta cat cta tac cag gat ctg 288
 Ala Leu Tyr Leu Gln Lys Thr Ala Asn Leu His Leu Tyr Gln Asp Leu
 85 90 95

caa aga gaa gtt ggg tct cta aaa gaa att aat ttc atg ctg agc gtt 336
 Gln Arg Glu Val Gly Ser Leu Lys Glu Ile Asn Phe Met Leu Ser Val
 100 105 110

cta cag aaa gaa ttt ctt cat tta tct aaa gaa ttt gca acg aca tct 384
 Leu Gln Lys Glu Phe Leu His Leu Ser Lys Glu Phe Ala Thr Thr Ser
 115 120 125

aaa gac ctc tct gct gta tct caa gat ttt tat tct tgt ttg caa gga 432
 Lys Asp Leu Ser Ala Val Ser Gln Asp Phe Tyr Ser Cys Leu Gln Gly
 130 135 140

ttt aga gat aac tat aaa ggt ttt gaa tct ctt ttg gat gag tat aaa 480
 Phe Arg Asp Asn Tyr Lys Gly Phe Glu Ser Leu Leu Asp Glu Tyr Lys
 145 150 155 160

aac tct aca gaa gaa atg cgc aaa ctc ttt tcg caa gaa atc ata gca 528


```

Asn Ser Thr Glu Glu Met Arg Lys Leu Phe Ser Gln Glu Ile Ile Ala
      165                      170                      175

gat ctt aaa ggc tct gtt gcc tca tta aga gag gaa atc cga ttc cta   576
Asp Leu Lys Gly Ser Val Ala Ser Leu Arg Glu Glu Ile Arg Phe Leu
      180                      185                      190

acc cca tta gca gaa gaa gtt cgc cga tta gcg cat aac cag gaa tca   624
Thr Pro Leu Ala Glu Glu Val Arg Arg Leu Ala His Asn Gln Glu Ser
      195                      200                      205

tta aca gcg gct att gaa gaa tta aaa aca att cgt gat agc tta cga   672
Leu Thr Ala Ala Ile Glu Glu Leu Lys Thr Ile Arg Asp Ser Leu Arg
      210                      215                      220

gat gaa att gga caa ctt tca caa ctt tct aaa act ctt acc agt caa   720
Asp Glu Ile Gly Gln Leu Ser Gln Leu Ser Lys Thr Leu Thr Ser Gln
      225                      230                      235                      240

att gca tta caa cga aaa gag agc tca gat ctg tgt tcc cag ata aga   768
Ile Ala Leu Gln Arg Lys Glu Ser Ser Asp Leu Cys Ser Gln Ile Arg
      245                      250                      255

gag acg ctc tcc tcc ccc aga aag tct gca tca ccc tct aca aaa agc   816
Glu Thr Leu Ser Ser Pro Arg Lys Ser Ala Ser Pro Ser Thr Lys Ser
      260                      265                      270

tcc tag   822
Ser

<210> 14
<211> 273
<212> PRT
<213> Chlamydia trachomatis

<400> 14
Met Thr Thr Pro Thr Leu Ile Val Ile Pro Pro Ser Pro Pro Ala Pro
  1                      5                      10                      15

Ser Tyr Ser Ala Asn Arg Val Pro Gln Pro Ser Leu Met Asp Lys Ile
      20                      25                      30

Lys Lys Ile Ala Ala Ile Ala Ser Leu Ile Leu Ile Gly Thr Ile Gly
      35                      40                      45

Phe Leu Ala Leu Leu Gly His Leu Val Gly Phe Leu Ile Ala Pro Gln
      50                      55                      60

Ile Thr Ile Val Leu Leu Ala Leu Phe Ile Thr Ser Leu Ala Gly Asn
      65                      70                      75                      80

Ala Leu Tyr Leu Gln Lys Thr Ala Asn Leu His Leu Tyr Gln Asp Leu
      85                      90                      95

Gln Arg Glu Val Gly Ser Leu Lys Glu Ile Asn Phe Met Leu Ser Val
      100                      105                      110

Leu Gln Lys Glu Phe Leu His Leu Ser Lys Glu Phe Ala Thr Thr Ser
      115                      120                      125

```

Lys Asp Leu Ser Ala Val Ser Gln Asp Phe Tyr Ser Cys Leu Gln Gly
 130 135 140
 Phe Arg Asp Asn Tyr Lys Gly Phe Glu Ser Leu Leu Asp Glu Tyr Lys
 145 150 155 160
 Asn Ser Thr Glu Glu Met Arg Lys Leu Phe Ser Gln Glu Ile Ile Ala
 165 170 175
 Asp Leu Lys Gly Ser Val Ala Ser Leu Arg Glu Glu Ile Arg Phe Leu
 180 185 190
 Thr Pro Leu Ala Glu Glu Val Arg Arg Leu Ala His Asn Gln Glu Ser
 195 200 205
 Leu Thr Ala Ala Ile Glu Glu Leu Lys Thr Ile Arg Asp Ser Leu Arg
 210 215 220
 Asp Glu Ile Gly Gln Leu Ser Gln Leu Ser Lys Thr Leu Thr Ser Gln
 225 230 235 240
 Ile Ala Leu Gln Arg Lys Glu Ser Ser Asp Leu Cys Ser Gln Ile Arg
 245 250 255
 Glu Thr Leu Ser Ser Pro Arg Lys Ser Ala Ser Pro Ser Thr Lys Ser
 260 265 270

Ser

<210> 15
 <211> 348
 <212> DNA
 <213> Chlamydia trachomatis

<220>
 <221> CDS
 <222> (1)..(348)

<400> 15
 atg gtt cat tct gta tac aat tca ttg gct cca gaa ggt ttt agc caa 48
 Met Val His Ser Val Tyr Asn Ser Leu Ala Pro Glu Gly Phe Ser Gln
 1 5 10 15
 gtc tct att caa ccc agt cag att cca acc agc aaa aaa gta atg att 96
 Val Ser Ile Gln Pro Ser Gln Ile Pro Thr Ser Lys Lys Val Met Ile
 20 25 30
 gcg ata atg act ctt ttt gca ctc aca gcc att gca gca ata gtc ctt 144
 Ala Ile Met Thr Leu Phe Ala Leu Thr Ala Ile Ala Ala Ile Val Leu
 35 40 45
 tcc atc gtt aca gtt tgt gga ggg ttt cct ttt ctt ctt gct gca ctt 192
 Ser Ile Val Thr Val Cys Gly Gly Phe Pro Phe Leu Leu Ala Ala Leu
 50 55 60
 aac acc gta act att ggt gca tgc gta tcc ttg ccg gta ttc act tgc 240
 Asn Thr Val Thr Ile Gly Ala Cys Val Ser Leu Pro Val Phe Thr Cys
 65 70 75 80
 ata gct aca acg tta tta ctt ctt tgt ctc cgt aat atc gaa ctc cta 288

```

Ile Ala Thr Thr Leu Leu Leu Leu Cys Leu Arg Asn Ile Glu Leu Leu
      85                      90                      95

gcc aga ccg caa gta ttt acc ctc tcc act caa ttc agc cca aca aaa   336
Ala Arg Pro Gln Val Phe Thr Leu Ser Thr Gln Phe Ser Pro Thr Lys
      100                      105                      110

cct caa gaa tag                                           348
Pro Gln Glu
      115

```

<210> 16
 <211> 115
 <212> PRT
 <213> Chlamydia trachomatis

```

<400> 16
Met Val His Ser Val Tyr Asn Ser Leu Ala Pro Glu Gly Phe Ser Gln
  1                      5                      10                      15

Val Ser Ile Gln Pro Ser Gln Ile Pro Thr Ser Lys Lys Val Met Ile
      20                      25                      30

Ala Ile Met Thr Leu Phe Ala Leu Thr Ala Ile Ala Ala Ile Val Leu
      35                      40                      45

Ser Ile Val Thr Val Cys Gly Gly Phe Pro Phe Leu Leu Ala Ala Leu
      50                      55                      60

Asn Thr Val Thr Ile Gly Ala Cys Val Ser Leu Pro Val Phe Thr Cys
      65                      70                      75                      80

Ile Ala Thr Thr Leu Leu Leu Leu Cys Leu Arg Asn Ile Glu Leu Leu
      85                      90                      95

Ala Arg Pro Gln Val Phe Thr Leu Ser Thr Gln Phe Ser Pro Thr Lys
      100                      105                      110

Pro Gln Glu
      115

```

<210> 17
 <211> 537
 <212> DNA
 <213> Chlamydia trachomatis

<220>
 <221> CDS
 <222> (1)..(537)

```

<400> 17
atg acg tac tct ata tcc gat ata gca cac aaa tct gat att tct aat   48
Met Thr Tyr Ser Ile Ser Asp Ile Ala His Lys Ser Asp Ile Ser Asn
  1                      5                      10                      15

ccc acg tct ccc gct cca tca aga aaa cga gga tcc ttt ccc cca caa   96
Pro Thr Ser Pro Ala Pro Ser Arg Lys Arg Gly Ser Phe Pro Pro Gln
      20                      25                      30

```

```

tct cct tct gcc gtg ggc tct tta gag gga gct aat ttc tct aca tgg 144
Ser Pro Ser Ala Val Gly Ser Leu Glu Gly Ala Asn Phe Ser Thr Trp
      35              40              45

ggg cca ggc ccc ttc ttc act gtc cct gtt tat cca caa caa ctc gct 192
Gly Pro Gly Pro Phe Phe Thr Val Pro Val Tyr Pro Gln Gln Leu Ala
      50              55              60

gca atg caa aac aac ctt ttt aca ttg caa aca gag gtt tct gct ctc 240
Ala Met Gln Asn Asn Leu Phe Thr Leu Gln Thr Glu Val Ser Ala Leu
      65              70              75              80

aag aaa aaa tta gtt cag tct agt cag aca cgc gga tct tta gga ctc 288
Lys Lys Lys Leu Val Gln Ser Ser Gln Thr Arg Gly Ser Leu Gly Leu
      85              90              95

ggc ccg cag ttt tta gcg gca tgc tta gtt gct gcg aca atc ctt gca 336
Gly Pro Gln Phe Leu Ala Ala Cys Leu Val Ala Ala Thr Ile Leu Ala
      100              105              110

gta gct gtt atc gta ctt gct tcc tta gga ctt ggc ggt gtt ctt cct 384
Val Ala Val Ile Val Leu Ala Ser Leu Gly Leu Gly Gly Val Leu Pro
      115              120              125

ttt gtc ctt gtt tgt ctg gct ggg tca act aat gca att tgg gct att 432
Phe Val Leu Val Cys Leu Ala Gly Ser Thr Asn Ala Ile Trp Ala Ile
      130              135              140

gtg agc gcc tcc atc act aca ctg att tgt tgc gtt tcc atc gct tgc 480
Val Ser Ala Ser Ile Thr Thr Leu Ile Cys Cys Val Ser Ile Ala Cys
      145              150              155              160

atc ttc tta gca aaa tgt gat aag gga tct gat cct caa act tta tat 528
Ile Phe Leu Ala Lys Cys Asp Lys Gly Ser Asp Pro Gln Thr Leu Tyr
      165              170              175

gta agc taa 537
Val Ser

```

<210> 18

<211> 178

<212> PRT

<213> Chlamydia trachomatis

<400> 18

```

Met Thr Tyr Ser Ile Ser Asp Ile Ala His Lys Ser Asp Ile Ser Asn
  1              5              10              15

```

```

Pro Thr Ser Pro Ala Pro Ser Arg Lys Arg Gly Ser Phe Pro Pro Gln
      20              25              30

```

```

Ser Pro Ser Ala Val Gly Ser Leu Glu Gly Ala Asn Phe Ser Thr Trp
      35              40              45

```

```

Gly Pro Gly Pro Phe Phe Thr Val Pro Val Tyr Pro Gln Gln Leu Ala
      50              55              60

```

```

Ala Met Gln Asn Asn Leu Phe Thr Leu Gln Thr Glu Val Ser Ala Leu
      65              70              75              80

```

Val Ser

22

30

23

19

<220>
<223> Description of Artificial Sequence: PCR primer

<400> 22
gcgcggatcc ttaattagtg ccttctggat tagg 34

<210> 23
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR primer

<400> 23
cgcagtactg tatccacaga caac 24

<210> 24
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR primer

<400> 24
gtcggatccg agaaactctc catgcc 26

WO 99/53948

PCT/US99/08744

SEQUENCE LISTING

<110> Oregon State University

<120> Methods of use for infection-specific INCA, INCB, and
INCC proteins of Chlamydia

<130> 52297

<140>

<141>

<150> 60/082,588

<151> 1998-04-21

<150> 60/082,438

<151> 1998-04-20

<150> 60/086,450

<151> 1998-05-22

<160> 24

<170> PatentIn Ver. 2.0

<210> 1

<211> 534

<212> DNA

<213> Chlamydia trachomatis

<220>

<221> CDS

<222> (1)..(534)

<400> 1

atg	aaa	aag	ttc	tta	tta	ctt	agc	tta	atg	tct	ttg	tca	tct	cta	cct	48
Met	Lys	Lys	Phe	Leu	Leu	Leu	Ser	Leu	Met	Ser	Leu	Ser	Ser	Leu	Pro	
1				5				10						15		

aca	ttt	gca	gct	aat	tct	aca	ggc	aca	att	gga	atc	gtt	aat	tta	cgt	96
Thr	Phe	Ala	Ala	Asn	Ser	Thr	Gly	Thr	Ile	Gly	Ile	Val	Asn	Leu	Arg	
			20					25					30			

cgc	tgc	cta	gaa	gag	tct	gct	ctt	ggg	aaa	aaa	gaa	tct	gct	gaa	ttc	144
Arg	Cys	Leu	Glu	Glu	Ser	Ala	Leu	Gly	Lys	Lys	Glu	Ser	Ala	Glu	Phe	
		35					40					45				

gaa	aag	atg	aaa	aac	caa	ttc	tct	aac	agc	atg	ggg	aag	atg	gag	gaa	192
Glu	Lys	Met	Lys	Asn	Gln	Phe	Ser	Asn	Ser	Met	Gly	Lys	Met	Glu	Glu	
	50					55					60					

gaa	ctg	tct	tct	atc	tat	tcc	aag	ctc	caa	gac	gac	gat	tac	atg	gaa	240
Glu	Leu	Ser	Ser	Ile	Tyr	Ser	Lys	Leu	Gln	Asp	Asp	Asp	Tyr	Met	Glu	
	65				70					75					80	

ggt	cta	tcc	gag	acc	gca	gct	gcc	gaa	tta	aga	aaa	aaa	ttc	gaa	gat	288
Gly	Leu	Ser	Glu	Thr	Ala	Ala	Ala	Glu	Leu	Arg	Lys	Lys	Phe	Glu	Asp	
			85					90					95			

cta	tct	gca	gaa	tac	aac	aca	gct	caa	ggg	cag	tat	tac	caa	ata	tta	336
Leu	Ser	Ala	Glu	Tyr	Asn	Thr	Ala	Gln	Gly	Gln	Tyr	Tyr	Gln	Ile	Leu	

100	105	110	
aac caa agt aat ttc aag cgc atg caa aag att atg gaa gaa gtg aaa			384
Asn Gln Ser Asn Phe Lys Arg Met Gln Lys Ile Met Glu Glu Val Lys			
115	120	125	
aaa gct tct gaa act gtg cgt att caa gaa ggc ttg tca gtc ctt ctt			432
Lys Ala Ser Glu Thr Val Arg Ile Gln Glu Gly Leu Ser Val Leu Leu			
130	135	140	
aac gaa gat att gtc tta tct atc gat agt tcg gca gat aaa acc gat			480
Asn Glu Asp Ile Val Leu Ser Ile Asp Ser Ser Ala Asp Lys Thr Asp			
145	150	155	160
gct gtt att aaa gtt ctt gat gtt ctt ttc aaa ata att aac atg cga			528
Ala Val Ile Lys Val Leu Asp Val Leu Phe Lys Ile Ile Asn Met Arg			
165	170	175	
agc tag			534
Ser			

<210> 2

<211> 177

<212> PRT

<213> Chlamydia trachomatis

<400> 2

Met Lys Lys Phe Leu Leu Leu Ser Leu Met Ser Leu Ser Ser Leu Pro			
1	5	10	15
Thr Phe Ala Ala Asn Ser Thr Gly Thr Ile Gly Ile Val Asn Leu Arg			
20	25	30	
Arg Cys Leu Glu Glu Ser Ala Leu Gly Lys Lys Glu Ser Ala Glu Phe			
35	40	45	
Glu Lys Met Lys Asn Gln Phe Ser Asn Ser Met Gly Lys Met Glu Glu			
50	55	60	
Glu Leu Ser Ser Ile Tyr Ser Lys Leu Gln Asp Asp Asp Tyr Met Glu			
65	70	75	80
Gly Leu Ser Glu Thr Ala Ala Ala Glu Leu Arg Lys Lys Phe Glu Asp			
85	90	95	
Leu Ser Ala Glu Tyr Asn Thr Ala Gln Gly Gln Tyr Tyr Gln Ile Leu			
100	105	110	
Asn Gln Ser Asn Phe Lys Arg Met Gln Lys Ile Met Glu Glu Val Lys			
115	120	125	
Lys Ala Ser Glu Thr Val Arg Ile Gln Glu Gly Leu Ser Val Leu Leu			
130	135	140	
Asn Glu Asp Ile Val Leu Ser Ile Asp Ser Ser Ala Asp Lys Thr Asp			
145	150	155	160
Ala Val Ile Lys Val Leu Asp Val Leu Phe Lys Ile Ile Asn Met Arg			
165	170	175	

Ser

<210> 3

<211> 846

<212> DNA

<213> Chlamydia trachomatis

<220>

<221> CDS

<222> (1)..(846)

<400> 3

```

atg aat cgc atg att tgt gat tgc gtg tct cgc ata act ggg gat cga      48
Met Asn Arg Met Ile Cys Asp Cys Val Ser Arg Ile Thr Gly Asp Arg
   1             5             10             15

gtc aag aat att gtt ctg att gat gga gcg att gat cct cat tca tat      96
Val Lys Asn Ile Val Leu Ile Asp Gly Ala Ile Asp Pro His Ser Tyr
           20             25             30

gag atg gtg aag ggg gat gaa gac cga atg gct atg agc cag ctg att      144
Glu Met Val Lys Gly Asp Glu Asp Arg Met Ala Met Ser Gln Leu Ile
           35             40             45

ttt tgc aat ggt tta ggt tta gag cat tca gct agt tta cgt aaa cat      192
Phe Cys Asn Gly Leu Gly Leu Glu His Ser Ala Ser Leu Arg Lys His
           50             55             60

cta gag ggt aac cca aaa gtc gtt gat tta ggt caa cgt ttg ctt aac      240
Leu Glu Gly Asn Pro Lys Val Val Asp Leu Gly Gln Arg Leu Leu Asn
           65             70             75             80

aaa aac tgt ttt gat ctt ctg agt gaa gaa gga ttc cct gac cca cat      288
Lys Asn Cys Phe Asp Leu Leu Ser Glu Glu Gly Phe Pro Asp Pro His
           85             90             95

att tgg acg gat atg aga gta tgg ggt gct gct gta aaa gag atg gct      336
Ile Trp Thr Asp Met Arg Val Trp Gly Ala Ala Val Lys Glu Met Ala
           100            105            110

gcg gca tta att caa caa ttt cct caa tat gaa gaa gat ttt caa aag      384
Ala Ala Leu Ile Gln Gln Phe Pro Gln Tyr Glu Glu Asp Phe Gln Lys
           115            120            125

aat gcg gat cag atc tta tca gag atg gag gaa ctt gat cgt tgg gca      432
Asn Ala Asp Gln Ile Leu Ser Glu Met Glu Glu Leu Asp Arg Trp Ala
           130            135            140

gtg cgt tct ctc tct acg att cct gaa aaa aat cgc tat tta gtc aca      480
Val Arg Ser Leu Ser Thr Ile Pro Glu Lys Asn Arg Tyr Leu Val Thr
           145            150            155            160

ggc cac aat gcg ttc agt tac ttt act cgt cgg tat cta tcc tct gat      528
Gly His Asn Ala Phe Ser Tyr Phe Thr Arg Arg Tyr Leu Ser Ser Asp
           165            170            175

gcg gag aga gtg tct ggg gaa tgg aga tcg cgt tgc att tct cca gaa      576
Ala Glu Arg Val Ser Gly Glu Trp Arg Ser Arg Cys Ile Ser Pro Glu
           180            185            190

```

ggg ttg tct cct gag gct cag att agt atc cga gat att atg cgt gta 624
 Gly Leu Ser Pro Glu Ala Gln Ile Ser Ile Arg Asp Ile Met Arg Val
 195 200 205
 gtg gag tat atc tct gca aac gat gta gaa gtt gtc ttt tta gag gat 672
 Val Glu Tyr Ile Ser Ala Asn Asp Val Glu Val Val Phe Leu Glu Asp
 210 215 220
 acg tta aat caa gat gct ttg aga aag att gtt tct tgc tct aag agc 720
 Thr Leu Asn Gln Asp Ala Leu Arg Lys Ile Val Ser Cys Ser Lys Ser
 225 230 235 240
 gga caa aag att cgt ctc gct aag tct cct tta tat agc gat aat gtc 768
 Gly Gln Lys Ile Arg Leu Ala Lys Ser Pro Leu Tyr Ser Asp Asn Val
 245 250 255
 tgt gat aac tat ttt agc acg ttc cag cac aat gtt cgc aca att aca 816
 Cys Asp Asn Tyr Phe Ser Thr Phe Gln His Asn Val Arg Thr Ile Thr
 260 265 270
 gaa gaa ttg gga ggg act gtt ctt gaa tag 846
 Glu Glu Leu Gly Gly Thr Val Leu Glu
 275 280

<210> 4

<211> 281

<212> PRT

<213> Chlamydia trachomatis

<400> 4

Met Asn Arg Met Ile Cys Asp Cys Val Ser Arg Ile Thr Gly Asp Arg
 1 5 10 15
 Val Lys Asn Ile Val Leu Ile Asp Gly Ala Ile Asp Pro His Ser Tyr
 20 25 30
 Glu Met Val Lys Gly Asp Glu Asp Arg Met Ala Met Ser Gln Leu Ile
 35 40 45
 Phe Cys Asn Gly Leu Gly Leu Glu His Ser Ala Ser Leu Arg Lys His
 50 55 60
 Leu Glu Gly Asn Pro Lys Val Val Asp Leu Gly Gln Arg Leu Leu Asn
 65 70 75 80
 Lys Asn Cys Phe Asp Leu Leu Ser Glu Glu Gly Phe Pro Asp Pro His
 85 90 95
 Ile Trp Thr Asp Met Arg Val Trp Gly Ala Ala Val Lys Glu Met Ala
 100 105 110
 Ala Ala Leu Ile Gln Gln Phe Pro Gln Tyr Glu Glu Asp Phe Gln Lys
 115 120 125
 Asn Ala Asp Gln Ile Leu Ser Glu Met Glu Glu Leu Asp Arg Trp Ala
 130 135 140
 Val Arg Ser Leu Ser Thr Ile Pro Glu Lys Asn Arg Tyr Leu Val Thr
 145 150 155 160

Gly His Asn Ala Phe Ser Tyr Phe Thr Arg Arg Tyr Leu Ser Ser Asp
 165 170 175

Ala Glu Arg Val Ser Gly Glu Trp Arg Ser Arg Cys Ile Ser Pro Glu
 180 185 190

Gly Leu Ser Pro Glu Ala Gln Ile Ser Ile Arg Asp Ile Met Arg Val
 195 200 205

Val Glu Tyr Ile Ser Ala Asn Asp Val Glu Val Val Phe Leu Glu Asp
 210 215 220

Thr Leu Asn Gln Asp Ala Leu Arg Lys Ile Val Ser Cys Ser Lys Ser
 225 230 235 240

Gly Gln Lys Ile Arg Leu Ala Lys Ser Pro Leu Tyr Ser Asp Asn Val
 245 250 255

Cys Asp Asn Tyr Phe Ser Thr Phe Gln His Asn Val Arg Thr Ile Thr
 260 265 270

Glu Glu Leu Gly Gly Thr Val Leu Glu
 275 280

<210> 5
 <211> 861
 <212> DNA
 <213> Chlamydia trachomatis

<220>
 <221> CDS
 <222> (1)..(861)

<400> 5
 atg tct gtg ata act att tta gca cgt tcc agc aca atg ttc gca caa 48
 Met Ser Val Ile Thr Ile Leu Ala Arg Ser Ser Thr Met Phe Ala Gln
 1 5 10 15

tta cag aag aat tgg gag gga ctg ttc ttg aat aga gat aat gca att 96
 Leu Gln Lys Asn Trp Glu Gly Leu Phe Leu Asn Arg Asp Asn Ala Ile
 20 25 30

gct tgg tcc gta gag gat ctt tgt gtt aat tat gat cac tca gac gtc 144
 Ala Trp Ser Val Glu Asp Leu Cys Val Asn Tyr Asp His Ser Asp Val
 35 40 45

tta tgt cac att act ttt tct ctg cct gca ggg gca atg gct gct att 192
 Leu Cys His Ile Thr Phe Ser Leu Pro Ala Gly Ala Met Ala Ala Ile
 50 55 60

att ggg ccg aat gga gct ggt aaa agt act ttg ctt aag gct tct tta 240
 Ile Gly Pro Asn Gly Ala Gly Lys Ser Thr Leu Leu Lys Ala Ser Leu
 65 70 75 80

gga ctg att cgt gct tct tct ggc caa agc ttg ttc ttt ggt cag aga 288
 Gly Leu Ile Arg Ala Ser Ser Gly Gln Ser Leu Phe Phe Gly Gln Arg
 85 90 95

ttt tcc aag gca cat cat aga ata gcc tat atg cct caa aga gcg agt 336
 Phe Ser Lys Ala His His Arg Ile Ala Tyr Met Pro Gln Arg Ala Ser

100	105	110	
gtg gat tgg gat ttc cca atg act gtt ctt gat ctc gtg ttg atg ggg Val Asp Trp Asp Phe Pro Met Thr Val Leu Asp Leu Val Leu Met Gly 115 120 125			384
tgt tac ggc tat aaa gga ata tgg aat cgt att tcc act gat gat cgt Cys Tyr Gly Tyr Lys Gly Ile Trp Asn Arg Ile Ser Thr Asp Asp Arg 130 135 140			432
cag gag gct atg cgt att tta gag cgg gtt ggt ttg gaa gct ttt gca Gln Glu Ala Met Arg Ile Leu Glu Arg Val Gly Leu Glu Ala Phe Ala 145 150 155 160			480
aat cgt caa ata ggt aag ctc tct gga gga caa caa cag aga gct ttt Asn Arg Gln Ile Gly Lys Leu Ser Gly Gln Gln Gln Arg Ala Phe 165 170 175			528
tta gcg cgg tca tta atg caa aaa gca gat ttg tat ctc atg gat gag Leu Ala Arg Ser Leu Met Gln Lys Ala Asp Leu Tyr Leu Met Asp Glu 180 185 190			576
ctg ttc tct gcg atc gat atg gcc tct tat cag atg gtt gta gat gtt Leu Phe Ser Ala Ile Asp Met Ala Ser Tyr Gln Met Val Val Asp Val 195 200 205			624
ttg caa gag ctt aaa agc gaa ggg aag act att gtg gtc att cat cat Leu Gln Glu Leu Lys Ser Glu Gly Lys Thr Ile Val Val Ile His His 210 215 220			672
gat ttg agt aat gtc cgg aag ctt ttt gat cat gtg att tta tta aat Asp Leu Ser Asn Val Arg Lys Leu Phe Asp His Val Ile Leu Leu Asn 225 230 235 240			720
aag cat ctt gtg tgc tct gga agc gta gaa gaa tgc ttg act aaa gaa Lys His Leu Val Cys Ser Gly Ser Val Glu Glu Cys Leu Thr Lys Glu 245 250 255			768
gcc att ttt cag gct tat ggg tgt gac ttg agc ttt tgg att aca cac Ala Ile Phe Gln Ala Tyr Gly Cys Asp Leu Ser Phe Trp Ile Thr His 260 265 270			816
tca aat tgt cta gag gca agt acc aag gat cgt gct aga tgc tga Ser Asn Cys Leu Glu Ala Ser Thr Lys Asp Arg Ala Arg Cys 275 280 285			861

<210> 6

<211> 286

<212> PRT

<213> Chlamydia trachomatis

<400> 6

Met	Ser	Val	Ile	Thr	Ile	Leu	Ala	Arg	Ser	Ser	Thr	Met	Phe	Ala	Gln
1					5					10				15	

Leu	Gln	Lys	Asn	Trp	Glu	Gly	Leu	Phe	Leu	Asn	Arg	Asp	Asn	Ala	Ile
		20					25						30		

Ala	Trp	Ser	Val	Glu	Asp	Leu	Cys	Val	Asn	Tyr	Asp	His	Ser	Asp	Val
		35					40					45			

Leu Cys His Ile Thr Phe Ser Leu Pro Ala Gly Ala Met Ala Ala Ile
 50 55 60
 Ile Gly Pro Asn Gly Ala Gly Lys Ser Thr Leu Leu Lys Ala Ser Leu
 65 70 75 80
 Gly Leu Ile Arg Ala Ser Ser Gly Gln Ser Leu Phe Phe Gly Gln Arg
 85 90 95
 Phe Ser Lys Ala His His Arg Ile Ala Tyr Met Pro Gln Arg Ala Ser
 100 105 110
 Val Asp Trp Asp Phe Pro Met Thr Val Leu Asp Leu Val Leu Met Gly
 115 120 125
 Cys Tyr Gly Tyr Lys Gly Ile Trp Asn Arg Ile Ser Thr Asp Asp Arg
 130 135 140
 Gln Glu Ala Met Arg Ile Leu Glu Arg Val Gly Leu Glu Ala Phe Ala
 145 150 155 160
 Asn Arg Gln Ile Gly Lys Leu Ser Gly Gly Gln Gln Gln Arg Ala Phe
 165 170 175
 Leu Ala Arg Ser Leu Met Gln Lys Ala Asp Leu Tyr Leu Met Asp Glu
 180 185 190
 Leu Phe Ser Ala Ile Asp Met Ala Ser Tyr Gln Met Val Val Asp Val
 195 200 205
 Leu Gln Glu Leu Lys Ser Glu Gly Lys Thr Ile Val Val Ile His His
 210 215 220
 Asp Leu Ser Asn Val Arg Lys Leu Phe Asp His Val Ile Leu Leu Asn
 225 230 235 240
 Lys His Leu Val Cys Ser Gly Ser Val Glu Glu Cys Leu Thr Lys Glu
 245 250 255
 Ala Ile Phe Gln Ala Tyr Gly Cys Asp Leu Ser Phe Trp Ile Thr His
 260 265 270
 Ser Asn Cys Leu Glu Ala Ser Thr Lys Asp Arg Ala Arg Cys
 275 280 285

<210> 7

<211> 1068

<212> DNA

<213> Chlamydia psittaci

<220>

<221> CDS

<222> (1)..(1068)

<400> 7

atg aca gta tcc aca gac aac aca agt cct gta ata tcg aga gcg tcc 48
 Met Thr Val Ser Thr Asp Asn Thr Ser Pro Val Ile Ser Arg Ala Ser
 1 5 10 15

tca cct act ttt gga gat cat ggt aag gat ttc gac aac aat aaa att	96
Ser Pro Thr Phe Gly Asp His Gly Lys Asp Phe Asp Asn Asn Lys Ile	
20 25 30	
ata ccc att tca ata gaa gct cca act tct tca gct gct gct gta ggg	144
Ile Pro Ile Ser Ile Glu Ala Pro Thr Ser Ser Ala Ala Ala Val Gly	
35 40 45	
gct aaa acg gct atc gag cct gaa gga aga agc cca cta ctt caa agg	192
Ala Lys Thr Ala Ile Glu Pro Glu Gly Arg Ser Pro Leu Leu Gln Arg	
50 55 60	
att tgc tat ctt gtt aaa att atc gct gcc atc gcc ctc ttt gtt gtt	240
Ile Cys Tyr Leu Val Lys Ile Ile Ala Ala Ile Ala Leu Phe Val Val	
65 70 75 80	
ggt atc gca gcc tta gtt tgc tta tat ctc ggt agc gtt atc tca acg	288
Gly Ile Ala Ala Leu Val Cys Leu Tyr Leu Gly Ser Val Ile Ser Thr	
85 90 95	
cct tct ctt att ctt atg ctt gcg atc atg ctt gta tcc ttt gtg atc	336
Pro Ser Leu Ile Leu Met Leu Ala Ile Met Leu Val Ser Phe Val Ile	
100 105 110	
gtt att acg gca att cga gat ggc aca ccg tct caa gtg gtc cgt cac	384
Val Ile Thr Ala Ile Arg Asp Gly Thr Pro Ser Gln Val Val Arg His	
115 120 125	
atg aaa cag caa att cag caa ttt ggc gaa gaa aac acg cgt tta cat	432
Met Lys Gln Gln Ile Gln Gln Phe Gly Glu Glu Asn Thr Arg Leu His	
130 135 140	
acc gca gta gaa aat cta aaa gct gtt aac gtt gag ctc tca gag caa	480
Thr Ala Val Glu Asn Leu Lys Ala Val Asn Val Glu Leu Ser Glu Gln	
145 150 155 160	
att aac caa ctt aaa caa cta cat act aga tta tcg gat ttt ggt gat	528
Ile Asn Gln Leu Lys Gln Leu His Thr Arg Leu Ser Asp Phe Gly Asp	
165 170 175	
agg ctt gaa gcg aat acc ggt gat ttt act gca ctt att gcg gat ttc	576
Arg Leu Glu Ala Asn Thr Gly Asp Phe Thr Ala Leu Ile Ala Asp Phe	
180 185 190	
caa ctc agt ctg gaa gag ttt aag tct gtt ggt act aaa gtt gaa acc	624
Gln Leu Ser Leu Glu Glu Phe Lys Ser Val Gly Thr Lys Val Glu Thr	
195 200 205	
atg ctc tct cca ttt gag aaa tta gct cag tct ttg aaa gag acc ttt	672
Met Leu Ser Pro Phe Glu Lys Leu Ala Gln Ser Leu Lys Glu Thr Phe	
210 215 220	
tct caa gaa gct gtt cag gca atg atg tcc tct gta act gag tta aga	720
Ser Gln Glu Ala Val Gln Ala Met Met Ser Ser Val Thr Glu Leu Arg	
225 230 235 240	
acc aat ttg aat gca ttg aaa gag ctt ata aca gag aat aaa acc gta	768
Thr Asn Leu Asn Ala Leu Lys Glu Leu Ile Thr Glu Asn Lys Thr Val	
245 250 255	
ata gag caa cta aaa gct gat gct caa ctt aga gaa gag caa gtg cgg	816

```

Ile Glu Gln Leu Lys Ala Asp Ala Gln Leu Arg Glu Glu Gln Val Arg
      260                      265                      270

ttt tta gaa aag cgt aaa caa gag tta gaa gag gct tgt tca aca ttg      864
Phe Leu Glu Lys Arg Lys Gln Glu Leu Glu Glu Ala Cys Ser Thr Leu
      275                      280                      285

tcc cat tca att gcg act cta cag gaa tcc aca acc ctt cta aag gac      912
Ser His Ser Ile Ala Thr Leu Gln Glu Ser Thr Thr Leu Leu Lys Asp
      290                      295                      300

tct aca act aac tta cat gca gtt gaa agt cgt ctt atc ggt gtt atg      960
Ser Thr Thr Asn Leu His Ala Val Glu Ser Arg Leu Ile Gly Val Met
      305                      310                      315                      320

gtt cag gat ggt gca gag tcc tcc acc gta gag gaa gct tca caa gat      1008
Val Gln Asp Gly Ala Glu Ser Ser Thr Val Glu Glu Ala Ser Gln Asp
      325                      330                      335

gat agc gcg caa ccc caa gat gaa aat caa tct gat gct gga gag cat      1056
Asp Ser Ala Gln Pro Gln Asp Glu Asn Gln Ser Asp Ala Gly Glu His
      340                      345                      350

aaa gat agt taa
Lys Asp Ser
      355

<210> 8
<211> 355
<212> PRT
<213> Chlamydia psittaci

<400> 8
Met Thr Val Ser Thr Asp Asn Thr Ser Pro Val Ile Ser Arg Ala Ser
  1                      5                      10                      15

Ser Pro Thr Phe Gly Asp His Gly Lys Asp Phe Asp Asn Asn Lys Ile
      20                      25                      30

Ile Pro Ile Ser Ile Glu Ala Pro Thr Ser Ser Ala Ala Ala Val Gly
      35                      40                      45

Ala Lys Thr Ala Ile Glu Pro Glu Gly Arg Ser Pro Leu Leu Gln Arg
      50                      55                      60

Ile Cys Tyr Leu Val Lys Ile Ile Ala Ala Ile Ala Leu Phe Val Val
      65                      70                      75                      80

Gly Ile Ala Ala Leu Val Cys Leu Tyr Leu Gly Ser Val Ile Ser Thr
      85                      90                      95

Pro Ser Leu Ile Leu Met Leu Ala Ile Met Leu Val Ser Phe Val Ile
      100                      105                      110

Val Ile Thr Ala Ile Arg Asp Gly Thr Pro Ser Gln Val Val Arg His
      115                      120                      125

Met Lys Gln Gln Ile Gln Gln Phe Gly Glu Glu Asn Thr Arg Leu His
      130                      135                      140

```

WO 99/53948

PCT/US99/08744

Thr Ala Val Glu Asn Leu Lys Ala Val Asn Val Glu Leu Ser Glu Gln
145 150 155 160

Ile Asn Gln Leu Lys Gln Leu His Thr Arg Leu Ser Asp Phe Gly Asp
165 170 175

Arg Leu Glu Ala Asn Thr Gly Asp Phe Thr Ala Leu Ile Ala Asp Phe
180 185 190

Gln Leu Ser Leu Glu Glu Phe Lys Ser Val Gly Thr Lys Val Glu Thr
195 200 205

Met Leu Ser Pro Phe Glu Lys Leu Ala Gln Ser Leu Lys Glu Thr Phe
210 215 220

Ser Gln Glu Ala Val Gln Ala Met Met Ser Ser Val Thr Glu Leu Arg
225 230 235 240

Thr Asn Leu Asn Ala Leu Lys Glu Leu Ile Thr Glu Asn Lys Thr Val
245 250 255

Ile Glu Gln Leu Lys Ala Asp Ala Gln Leu Arg Glu Glu Gln Val Arg
260 265 270

Phe Leu Glu Lys Arg Lys Gln Glu Leu Glu Glu Ala Cys Ser Thr Leu
275 280 285

Ser His Ser Ile Ala Thr Leu Gln Glu Ser Thr Thr Leu Leu Lys Asp
290 295 300

Ser Thr Thr Asn Leu His Ala Val Glu Ser Arg Leu Ile Gly Val Met
305 310 315 320

Val Gln Asp Gly Ala Glu Ser Ser Thr Val Glu Glu Ala Ser Gln Asp
325 330 335

Asp Ser Ala Gln Pro Gln Asp Glu Asn Gln Ser Asp Ala Gly Glu His
340 345 350

Lys Asp Ser
355

<210> 9
<211> 597
<212> DNA
<213> Chlamydia psittaci

<220>
<221> CDS
<222> (1)..(597)

<400> 9
atg tca aca aca cca gca tct tca gca agt cga gac gta tta tta gat 48
Met Ser Thr Thr Pro Ala Ser Ser Ala Ser Arg Asp Val Leu Leu Asp
1 5 10 15

gac gtt tta ata gct ttt aat aga aag cta aat ctc gta gaa caa caa 96
Asp Val Leu Ile Ala Phe Asn Arg Lys Leu Asn Leu Val Glu Gln Gln
20 25 30

gcg aaa gaa ctt gaa acg aaa gtc agt ttg gta gac aga aca gct act 144
Ala Lys Glu Leu Glu Thr Lys Val Ser Leu Val Asp Arg Thr Ala Thr
35 40 45

tta tca ctt acc act ggc aat aat gta gcc acg gat gta ctc ctt tta 192
Leu Ser Leu Thr Thr Gly Asn Asn Val Ala Thr Asp Val Leu Leu Leu
50 55 60

aaa gat gag gtt gca gaa cta aaa gga tgt ttg tct gca gtt acg gat 240
Lys Asp Glu Val Ala Glu Leu Lys Gly Cys Leu Ser Ala Val Thr Asp
65 70 75 80

cta tta atc cgc tca ggc tca tca aga aca cct ggg ggt gct cct aat 288
Leu Leu Ile Arg Ser Gly Ser Ser Arg Thr Pro Gly Gly Ala Pro Asn
85 90 95

cca gaa ggc act aat tac cta ata gga tgc aca cct cct tct ctt tgc 336
Pro Glu Gly Thr Asn Tyr Leu Ile Gly Cys Thr Pro Pro Ser Leu Cys
100 105 110

gct aaa ctt aca gcg tta gcg tta aca att ata gcc ctc att gct atc 384
Ala Lys Leu Thr Ala Leu Ala Leu Thr Ile Ile Ala Leu Ile Ala Ile
115 120 125

aca gta ctt gtt atc tgt att gtt act gtt tgc ggc ggt ttc ccc cta 432
Thr Val Leu Val Ile Cys Ile Val Thr Val Cys Gly Gly Phe Pro Leu
130 135 140

ttt att tcc cta ctc aac atg tac aca gtt ggt gct tgt ata tcc tta 480
Phe Ile Ser Leu Leu Asn Met Tyr Thr Val Gly Ala Cys Ile Ser Leu
145 150 155 160

ccg atc att tcg tgt gcc gca gtt tca atg atg att cta tgc tca cat 528
Pro Ile Ile Ser Cys Ala Ala Val Ser Met Met Ile Leu Cys Ser His
165 170 175

tct att aac tct tta tta aga aac agg cct gcg atc tat atg act aac 576
Ser Ile Asn Ser Leu Leu Arg Asn Arg Pro Ala Ile Tyr Met Thr Asn
180 185 190

aat ttt caa aca gaa tct taa 597
Asn Phe Gln Thr Glu Ser
195

<210> 10

<211> 198

<212> PRT

<213> Chlamydia psittaci

<400> 10

Met Ser Thr Thr Pro Ala Ser Ser Ala Ser Arg Asp Val Leu Leu Asp
1 5 10 15

Asp Val Leu Ile Ala Phe Asn Arg Lys Leu Asn Leu Val Glu Gln Gln
20 25 30

Ala Lys Glu Leu Glu Thr Lys Val Ser Leu Val Asp Arg Thr Ala Thr
35 40 45

Leu Ser Leu Thr Thr Gly Asn Asn Val Ala Thr Asp Val Leu Leu Leu

50 55 60
 Lys Asp Glu Val Ala Glu Leu Lys Gly Cys Leu Ser Ala Val Thr Asp
 65 70 75 80
 Leu Leu Ile Arg Ser Gly Ser Ser Arg Thr Pro Gly Gly Ala Pro Asn
 85 90 95
 Pro Glu Gly Thr Asn Tyr Leu Ile Gly Cys Thr Pro Pro Ser Leu Cys
 100 105 110
 Ala Lys Leu Thr Ala Leu Ala Leu Thr Ile Ile Ala Leu Ile Ala Ile
 115 120 125
 Thr Val Leu Val Ile Cys Ile Val Thr Val Cys Gly Gly Phe Pro Leu
 130 135 140
 Phe Ile Ser Leu Leu Asn Met Tyr Thr Val Gly Ala Cys Ile Ser Leu
 145 150 155 160
 Pro Ile Ile Ser Cys Ala Ala Val Ser Met Met Ile Leu Cys Ser His
 165 170 175
 Ser Ile Asn Ser Leu Leu Arg Asn Arg Pro Ala Ile Tyr Met Thr Asn
 180 185 190
 Asn Phe Gln Thr Glu Ser
 195

<210> 11
 <211> 561
 <212> DNA
 <213> Chlamydia psittaci

<220>
 <221> CDS
 <222> (1)..(561)

<400> 11
 atg acc tct gta aga acc gat tta act cca ggc gac acc tca ctc caa 48
 Met Thr Ser Val Arg Thr Asp Leu Thr Pro Gly Asp Thr Ser Leu Gln
 1 5 10 15
 tct tct tta tta aat ccg agt gat ctc aca aca caa cta tcc aac ctc 96
 Ser Ser Leu Leu Asn Pro Ser Asp Leu Thr Thr Gln Leu Ser Asn Leu
 20 25 30
 cag act gtt ctc gca ggg ata caa caa caa cat cct tta aac ggt ggt 144
 Gln Thr Val Leu Ala Gly Ile Gln Gln Gln His Pro Leu Asn Gly Gly
 35 40 45
 tgg cct cag cat cat cct act ggc gct gca gat caa aat tat ctc atg 192
 Trp Pro Gln His His Pro Thr Gly Ala Ala Asp Gln Asn Tyr Leu Met
 50 55 60
 cgt ctg atg caa tct cat atg gca agt acc gta tca gca gta tct gaa 240
 Arg Leu Met Gln Ser His Met Ala Ser Thr Val Ser Ala Val Ser Glu
 65 70 75 80
 tta aga acc gaa gtc act gca atc aag aca aaa ttg cac ggg cta tct 288

Leu Arg Thr Glu Val Thr Ala Ile Lys Thr Lys Leu His Gly Leu Ser
 85 90 95

 act cca gct aat gtt tgc agc ggt cct atg gct cta gcc gct ttt ctt 336
 Thr Pro Ala Asn Val Cys Ser Gly Pro Met Ala Leu Ala Ala Phe Leu
 100 105 110

 cta gct ata tct tta gtt gcg att atc atc att gtt tta gcc tcc tta 384
 Leu Ala Ile Ser Leu Val Ala Ile Ile Ile Ile Val Leu Ala Ser Leu
 115 120 125

 ggc ctt gca ggc ata cta cct caa gct gcc gct atc tta gtg aat aca 432
 Gly Leu Ala Gly Ile Leu Pro Gln Ala Ala Ala Ile Leu Val Asn Thr
 130 135 140

 gca aac tct ata tgg gct att gtt agc gct tcg ata gtc act gtt atc 480
 Ala Asn Ser Ile Trp Ala Ile Val Ser Ala Ser Ile Val Thr Val Ile
 145 150 155 160

 tgc tta att agc gtg cta tgc ata acg cta att cga cac cat aaa ccc 528
 Cys Leu Ile Ser Val Leu Cys Ile Thr Leu Ile Arg His His Lys Pro
 165 170 175

 tta cct att gaa act agg cct acc gga cat taa 561
 Leu Pro Ile Glu Thr Arg Pro Thr Gly His
 180 185

 <210> 12
 <211> 186
 <212> PRT
 <213> Chlamydia psittaci

 <400> 12
 Met Thr Ser Val Arg Thr Asp Leu Thr Pro Gly Asp Thr Ser Leu Gln
 1 5 10 15

 Ser Ser Leu Leu Asn Pro Ser Asp Leu Thr Thr Gln Leu Ser Asn Leu
 20 25 30

 Gln Thr Val Leu Ala Gly Ile Gln Gln Gln His Pro Leu Asn Gly Gly
 35 40 45

 Trp Pro Gln His His Pro Thr Gly Ala Ala Asp Gln Asn Tyr Leu Met
 50 55 60

 Arg Leu Met Gln Ser His Met Ala Ser Thr Val Ser Ala Val Ser Glu
 65 70 75 80

 Leu Arg Thr Glu Val Thr Ala Ile Lys Thr Lys Leu His Gly Leu Ser
 85 90 95

 Thr Pro Ala Asn Val Cys Ser Gly Pro Met Ala Leu Ala Ala Phe Leu
 100 105 110

 Leu Ala Ile Ser Leu Val Ala Ile Ile Ile Ile Val Leu Ala Ser Leu
 115 120 125

 Gly Leu Ala Gly Ile Leu Pro Gln Ala Ala Ala Ile Leu Val Asn Thr
 130 135 140

Ala Asn Ser Ile Trp Ala Ile Val Ser Ala Ser Ile Val Thr Val Ile
 145 150 155 160

Cys Leu Ile Ser Val Leu Cys Ile Thr Leu Ile Arg His His Lys Pro
 165 170 175

Leu Pro Ile Glu Thr Arg Pro Thr Gly His
 180 185

<210> 13

<211> 822

<212> DNA

<213> Chlamydia trachomatis

<220>

<221> CDS

<222> (1)..(822)

<400> 13

atg aca acg cct act cta atc gtg att cct cca tct ccc cct gca cct 48
 Met Thr Thr Pro Thr Leu Ile Val Ile Pro Pro Ser Pro Pro Ala Pro
 1 5 10 15

tcc tac tca gcc aat cgc gta cct caa cct tct ttg atg gac aaa att 96
 Ser Tyr Ser Ala Asn Arg Val Pro Gln Pro Ser Leu Met Asp Lys Ile
 20 25 30

aag aaa ata gca gcc att gcc tcc cta att ctt ata ggc aca ata ggc 144
 Lys Lys Ile Ala Ala Ile Ala Ser Leu Ile Leu Ile Gly Thr Ile Gly
 35 40 45

ttt tta gct ctt ttg gga cat ctt gtt ggc ttt ctg atc gct cca caa 192
 Phe Leu Ala Leu Leu Gly His Leu Val Gly Phe Leu Ile Ala Pro Gln
 50 55 60

atc act att gtt ctt ctt gcc cta ttc att acc tca tta gca ggc aat 240
 Ile Thr Ile Val Leu Leu Ala Leu Phe Ile Thr Ser Leu Ala Gly Asn
 65 70 75 80

gct ctt tat cta cag aaa acc gct aat cta cat cta tac cag gat ctg 288
 Ala Leu Tyr Leu Gln Lys Thr Ala Asn Leu His Leu Tyr Gln Asp Leu
 85 90 95

caa aga gaa gtt ggg tct cta aaa gaa att aat ttc atg ctg agc gtt 336
 Gln Arg Glu Val Gly Ser Leu Lys Glu Ile Asn Phe Met Leu Ser Val
 100 105 110

cta cag aaa gaa ttt ctt cat tta tct aaa gaa ttt gca acg aca tct 384
 Leu Gln Lys Glu Phe Leu His Leu Ser Lys Glu Phe Ala Thr Thr Ser
 115 120 125

aaa gac ctc tct gct gta tct caa gat ttt tat tct tgt ttg caa gga 432
 Lys Asp Leu Ser Ala Val Ser Gln Asp Phe Tyr Ser Cys Leu Gln Gly
 130 135 140

ttt aga gat aac tat aaa ggt ttt gaa tct ctt ttg gat gag tat aaa 480
 Phe Arg Asp Asn Tyr Lys Gly Phe Glu Ser Leu Leu Asp Glu Tyr Lys
 145 150 155 160

aac tct aca gaa gaa atg cgc aaa ctc ttt tcg caa gaa atc ata gca 528

```

Asn Ser Thr Glu Glu Met Arg Lys Leu Phe Ser Gln Glu Ile Ile Ala
      165                      170                      175

gat ctt aaa ggc tct gtt gcc tca tta aga gag gaa atc cga ttc cta   576
Asp Leu Lys Gly Ser Val Ala Ser Leu Arg Glu Glu Ile Arg Phe Leu
      180                      185                      190

acc cca tta gca gaa gaa gtt cgc cga tta gcg cat aac cag gaa tca   624
Thr Pro Leu Ala Glu Glu Val Arg Arg Leu Ala His Asn Gln Glu Ser
      195                      200                      205

tta aca gcg gct att gaa gaa tta aaa aca att cgt gat agc tta cga   672
Leu Thr Ala Ala Ile Glu Glu Leu Lys Thr Ile Arg Asp Ser Leu Arg
      210                      215                      220

gat gaa att gga caa ctt tca caa ctt tct aaa act ctt acc agt caa   720
Asp Glu Ile Gly Gln Leu Ser Gln Leu Ser Lys Thr Leu Thr Ser Gln
      225                      230                      235                      240

att gca tta caa cga aaa gag agc tca gat ctg tgt tcc cag ata aga   768
Ile Ala Leu Gln Arg Lys Glu Ser Ser Asp Leu Cys Ser Gln Ile Arg
      245                      250                      255

gag acg ctc tcc tcc ccc aga aag tct gca tca ccc tct aca aaa agc   816
Glu Thr Leu Ser Ser Pro Arg Lys Ser Ala Ser Pro Ser Thr Lys Ser
      260                      265                      270

tcc tag
Ser
      822

<210> 14
<211> 273
<212> PRT
<213> Chlamydia trachomatis

<400> 14
Met Thr Thr Pro Thr Leu Ile Val Ile Pro Pro Ser Pro Pro Ala Pro
  1                      5                      10                      15

Ser Tyr Ser Ala Asn Arg Val Pro Gln Pro Ser Leu Met Asp Lys Ile
      20                      25                      30

Lys Lys Ile Ala Ala Ile Ala Ser Leu Ile Leu Ile Gly Thr Ile Gly
      35                      40                      45

Phe Leu Ala Leu Leu Gly His Leu Val Gly Phe Leu Ile Ala Pro Gln
      50                      55                      60

Ile Thr Ile Val Leu Leu Ala Leu Phe Ile Thr Ser Leu Ala Gly Asn
      65                      70                      75                      80

Ala Leu Tyr Leu Gln Lys Thr Ala Asn Leu His Leu Tyr Gln Asp Leu
      85                      90                      95

Gln Arg Glu Val Gly Ser Leu Lys Glu Ile Asn Phe Met Leu Ser Val
      100                      105                      110

Leu Gln Lys Glu Phe Leu His Leu Ser Lys Glu Phe Ala Thr Thr Ser
      115                      120                      125

```

Lys Asp Leu Ser Ala Val Ser Gln Asp Phe Tyr Ser Cys Leu Gln Gly
 130 135 140
 Phe Arg Asp Asn Tyr Lys Gly Phe Glu Ser Leu Leu Asp Glu Tyr Lys
 145 150 155 160
 Asn Ser Thr Glu Glu Met Arg Lys Leu Phe Ser Gln Glu Ile Ile Ala
 165 170 175
 Asp Leu Lys Gly Ser Val Ala Ser Leu Arg Glu Glu Ile Arg Phe Leu
 180 185 190
 Thr Pro Leu Ala Glu Glu Val Arg Arg Leu Ala His Asn Gln Glu Ser
 195 200 205
 Leu Thr Ala Ala Ile Glu Glu Leu Lys Thr Ile Arg Asp Ser Leu Arg
 210 215 220
 Asp Glu Ile Gly Gln Leu Ser Gln Leu Ser Lys Thr Leu Thr Ser Gln
 225 230 235 240
 Ile Ala Leu Gln Arg Lys Glu Ser Ser Asp Leu Cys Ser Gln Ile Arg
 245 250 255
 Glu Thr Leu Ser Ser Pro Arg Lys Ser Ala Ser Pro Ser Thr Lys Ser
 260 265 270

Ser

<210> 15

<211> 348

<212> DNA

<213> Chlamydia trachomatis

<220>

<221> CDS

<222> (1)..(348)

<400> 15

atg gtt cat tct gta tac aat tca ttg gct cca gaa ggt ttt agc caa 48
 Met Val His Ser Val Tyr Asn Ser Leu Ala Pro Glu Gly Phe Ser Gln
 1 5 10 15

gtc tct att caa ccc agt cag att cca acc agc aaa aaa gta atg att 96
 Val Ser Ile Gln Pro Ser Gln Ile Pro Thr Ser Lys Lys Val Met Ile
 20 25 30

gcg ata atg act ctt ttt gca ctc aca gcc att gca gca ata gtc ctt 144
 Ala Ile Met Thr Leu Phe Ala Leu Thr Ala Ile Ala Ala Ile Val Leu
 35 40 45

tcc atc gtt aca gtt tgt gga ggg ttt cct ttt ctt ctt gct gca ctt 192
 Ser Ile Val Thr Val Cys Gly Gly Phe Pro Phe Leu Leu Ala Ala Leu
 50 55 60

aac acc gta act att ggt gca tgc gta tcc ttg ccg gta ttc act tgc 240
 Asn Thr Val Thr Ile Gly Ala Cys Val Ser Leu Pro Val Phe Thr Cys
 65 70 75 80

ata gct aca acg tta tta ctt ctt tgt ctc cgt aat atc gaa ctc cta 288

Ile Ala Thr Thr Leu Leu Leu Cys Leu Arg Asn Ile Glu Leu Leu
 85 90 95

gcc aga ccg caa gta ttt acc ctc tcc act caa ttc agc cca aca aaa 336
 Ala Arg Pro Gln Val Phe Thr Leu Ser Thr Gln Phe Ser Pro Thr Lys
 100 105 110

cct caa gaa tag 348
 Pro Gln Glu
 115

<210> 16
 <211> 115
 <212> PRT
 <213> Chlamydia trachomatis

<400> 16
 Met Val His Ser Val Tyr Asn Ser Leu Ala Pro Glu Gly Phe Ser Gln
 1 5 10 15

Val Ser Ile Gln Pro Ser Gln Ile Pro Thr Ser Lys Lys Val Met Ile
 20 25 30

Ala Ile Met Thr Leu Phe Ala Leu Thr Ala Ile Ala Ala Ile Val Leu
 35 40 45

Ser Ile Val Thr Val Cys Gly Gly Phe Pro Phe Leu Leu Ala Ala Leu
 50 55 60

Asn Thr Val Thr Ile Gly Ala Cys Val Ser Leu Pro Val Phe Thr Cys
 65 70 75 80

Ile Ala Thr Thr Leu Leu Leu Leu Cys Leu Arg Asn Ile Glu Leu Leu
 85 90 95

Ala Arg Pro Gln Val Phe Thr Leu Ser Thr Gln Phe Ser Pro Thr Lys
 100 105 110

Pro Gln Glu
 115

<210> 17
 <211> 537
 <212> DNA
 <213> Chlamydia trachomatis

<220>
 <221> CDS
 <222> (1)..(537)

<400> 17
 atg acg tac tct ata tcc gat ata gca cac aaa tct gat att tct aat 48
 Met Thr Tyr Ser Ile Ser Asp Ile Ala His Lys Ser Asp Ile Ser Asn
 1 5 10 15

ccc acg tct ccc gct cca tca aga aaa cga gga tcc ttt ccc cca caa 96
 Pro Thr Ser Pro Ala Pro Ser Arg Lys Arg Gly Ser Phe Pro Pro Gln
 20 25 30

tct cct tct gcc gtg ggc tct tta gag gga gct aat ttc tct aca tgg 144
 Ser Pro Ser Ala Val Gly Ser Leu Glu Gly Ala Asn Phe Ser Thr Trp
 35 40 45

ggg cca ggc ccc ttc ttc act gtc cct gtt tat cca caa caa ctc gct 192
 Gly Pro Gly Pro Phe Phe Thr Val Pro Val Tyr Pro Gln Gln Leu Ala
 50 55 60

gca atg caa aac aac ctt ttt aca ttg caa aca gag gtt tct gct ctc 240
 Ala Met Gln Asn Asn Leu Phe Thr Leu Gln Thr Glu Val Ser Ala Leu
 65 70 75 80

aag aaa aaa tta gtt cag tct agt cag aca cgc gga tct tta gga ctc 288
 Lys Lys Lys Leu Val Gln Ser Ser Gln Thr Arg Gly Ser Leu Gly Leu
 85 90 95

ggc ccg cag ttt tta gcg gca tgc tta gtt gct gcg aca atc ctt gca 336
 Gly Pro Gln Phe Leu Ala Ala Cys Leu Val Ala Ala Thr Ile Leu Ala
 100 105 110

gta gct gtt atc gta ctt gct tcc tta gga ctt ggc ggt gtt ctt cct 384
 Val Ala Val Ile Val Leu Ala Ser Leu Gly Leu Gly Gly Val Leu Pro
 115 120 125

ttt gtc ctt gtt tgt ctg gct ggg tca act aat gca att tgg gct att 432
 Phe Val Leu Val Cys Leu Ala Gly Ser Thr Asn Ala Ile Trp Ala Ile
 130 135 140

gtg agc gcc tcc atc act aca ctg att tgt tgc gtt tcc atc gct tgc 480
 Val Ser Ala Ser Ile Thr Thr Leu Ile Cys Cys Val Ser Ile Ala Cys
 145 150 155 160

atc ttc tta gca aaa tgt gat aag gga tct gat cct caa act tta tat 528
 Ile Phe Leu Ala Lys Cys Asp Lys Gly Ser Asp Pro Gln Thr Leu Tyr
 165 170 175

gta agc taa 537
 Val Ser

<210> 18

<211> 178

<212> PRT

<213> Chlamydia trachomatis

<400> 18

Met Thr Tyr Ser Ile Ser Asp Ile Ala His Lys Ser Asp Ile Ser Asn
 1 5 10 15

Pro Thr Ser Pro Ala Pro Ser Arg Lys Arg Gly Ser Phe Pro Pro Gln
 20 25 30

Ser Pro Ser Ala Val Gly Ser Leu Glu Gly Ala Asn Phe Ser Thr Trp
 35 40 45

Gly Pro Gly Pro Phe Phe Thr Val Pro Val Tyr Pro Gln Gln Leu Ala
 50 55 60

Ala Met Gln Asn Asn Leu Phe Thr Leu Gln Thr Glu Val Ser Ala Leu
 65 70 75 80

Lys Lys Lys Leu Val Gln Ser Ser Gln Thr Arg Gly Ser Leu Gly Leu
85 90 95

Gly Pro Gln Phe Leu Ala Ala Cys Leu Val Ala Ala Thr Ile Leu Ala
100 105 110

Val Ala Val Ile Val Leu Ala Ser Leu Gly Leu Gly Gly Val Leu Pro
115 120 125

Phe Val Leu Val Cys Leu Ala Gly Ser Thr Asn Ala Ile Trp Ala Ile
130 135 140

Val Ser Ala Ser Ile Thr Thr Leu Ile Cys Cys Val Ser Ile Ala Cys
145 150 155 160

Ile Phe Leu Ala Lys Cys Asp Lys Gly Ser Asp Pro Gln Thr Leu Tyr
165 170 175

Val Ser

<210> 19

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PCR primer

<400> 19

agaaccgatt taactccagg cg

22

<210> 20

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PCR primer

<400> 20

gcgcggatcc ttaatgtccg gtaggcctag

30

<210> 21

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PCR primer

<400> 21

atgtcaacaa caccagcatc ttc

23

<210> 22

<211> 34

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PCR primer

<400> 22

gcgcggatcc ttaattagtg ctttctggat tagg

34

<210> 23

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PCR primer

<400> 23

cgcagtactg tatccacaga caac

24

<210> 24

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PCR primer

<400> 24

gtcggatccg agaaactctc catgcc

26

PCT09

RAW SEQUENCE LISTING Date 01/18/2001
 PATENT APPLICATION NO. US/09/673,763 Title 16 34 38

Input Seq A:\55928 seq.txt
 Output Seq N:\CRF3\01182001\I673763.raw

ENTERED

9 <110> APPLICANT Oregon State University
 5 <120> TITLE OF INVENTION Methods of use for infection-specific INCA, INCB, and
 6 INCC proteins of Chlamydia
 8 <130> FILE REFERENCE 52257
 C--> 10 <140> CURRENT APPLICATION NUMBER: US/09/673,763
 C--> 11 <141> CURRENT FILING DATE: 2000-10-16
 13 <150> PRIOR APPLICATION NUMBER 60/082,566
 14 <151> PRIOR FILING DATE 1998-04-21
 16 <150> PRIOR APPLICATION NUMBER 60/082,433
 17 <151> PRIOR FILING DATE 1998-04-20
 19 <150> PRIOR APPLICATION NUMBER 60/086,156
 20 <151> PRIOR FILING DATE 1998-03-22
 22 <160> NUMBER OF SEQ ID NOS 24
 24 <170> SOFTWARE PatentIn Ver 2.0
 26 <210> SEQ ID NO 1
 27 <211> LENGTH 534
 28 <212> TYPE DNA
 29 <213> ORGANISM Chlamydia trachomatis
 31 <220> FEATURE
 32 <221> NAME/KEY CDS
 33 <222> LOCATION (1) (534)
 35 <400> SEQUENCE 1
 36 atg aag aag ttc tta tta ctt agc tta atg tct ttg tca tct cta cct 6
 37 Met Lys Lys Phe Leu Leu Leu Ser Lys Met Ser Leu Ser Ser Lys Pro
 38 1 5 10 15
 40 acc ttt gca gct aat tct tca ggc tca att ggc atc gtt tat tta cgt 96
 41 Thr Phe Ala Ala Asn Ser Thr Gly Thr Ala Gly Ile Val Asn Leu Arg
 42 20 25 30
 44 ggc tgc tta gaa gag tct gct ctt gag taa aat gca tct gct gaa ttt 144
 45 Arg Cys Leu Glu Glu Ser Ala Ieu Gly Lys Cys Glu Ser Ala Glu Phe
 46 35 40 45
 48 gaa aag atg aat aac caa ttc tct aac agc atg ggc aag atg gac gaa 192
 49 Glu Lys Met Lys Asn Gln Phe Ser Asn Ser Met Gly Lys Met Gly Glu
 50 50 55 60
 52 gaa atg tct tct atc tat tcc aag ctc caa gac gac gat tac atg gaa 240
 53 Glu Ieu Ser Ser Ile Tyr Ser Lys Ieu Gln Asp Asp Asp Tyr Met Glu
 54 65 70 75 80
 56 ggt cta tcc gag acc gca gcc gcc gaa tta aga aac aaa ttc gaa gtt 288
 57 Gly Leu Ser Glu Thr Ala Ala Ala Glu Leu Arg Phe Lys Phe Glu Asp
 58 85 90 95
 60 cta tct gca gaa tac aac caa gct cta ggc cag tct tac caa a a tta 336
 61 Leu Ser Ala Glu Tyr Asn Thr Ala Gln Gly Gln Tyr Tyr Gln Ile Leu
 62 100 105 110
 64 aac caa ggt aat ttc atg tgc atg caa aag ttt atg gaa gaa gta aat 384
 65 Asn Cln Ser Asn Phe Lys Arg Met Cln Lys Tie Met Glu Glu Val Lys
 66 115 120 125
 68 aaa gct tct gaa act gtg cct att caa gaa ggc ttg tca gtc ctt ctt 432

RAW SEQUENCE LISTING DATE: 01/18/2001
 PATENT APPLICATION US/09/673,763 TITLE: 1 2 3 38

Input Set A:\55928 seq.txt
 Output Set N:\CRF3\01182001\I673763.raw

```

69 Lys Ala Ser Ile Ile Val Arg Ile Gln Glu Gly Leu Ser Val Leu Leu
70      30      135      140
72 aac gaa gat att gta tta tct atc gat agt tgg gca cat aaa acc gat      480
73 Asn Glu asp ile val leu ser ile asp ser ser ala asp lys phe asp
74 145      150      155      160
76 gct att att aac gtt ctt gat gtt ctt ttc aaa ata att aac atg cga      528
77 Ala Val ile lys val leu aso val leu phe lys ile ile asn del val
78      165      170      175
80 aac tag      531
81 Ser
84 <210> SEQ ID NO: 2
85 <210> LENGTH: 177
86 <212> TYPE: PRT
87 <213> ORGANISM: Chlamydia trachomatis
89 <400> SEQUENCE:
90 Met Lys Lys Phe Leu Leu Leu Ser Ile Met Ser Leu Ser Ser Ile Pro
91      1      5      10      15
93 Thr Phe Ala Ala Asn ser thr Gly Thr Ile Glu Ile Val Asn Leu Arg
94      20      25      30
96 Arg Cys Leu Glu Glu Ser Ala Leu Glu Lys Lys Glu Ser Ala Glu Phe
97      35      40      45
99 Glu Lys Met Lys Phe Gln Phe Ser Asp Ser Met Gly Lys Met Glu Glu
100     50     55     60
102 Glu Leu Ser Ser Ile Thr Ser Lys Leu Cys Asp Asp Asp Tyr Met Glu
103     65     70     75     80
105 Glu Leu Ser Glu Thr Ala Ala Ala Glu Leu Arg Lys Lys Phe Glu Asp
106     85     90     95
108 Leu Ser Ala Glu Tyr Asn Thr Ala Cys Gly Gln Thr Thr Gln Ile Leu
109    100    105    110
111 Asn Gln Ser Asp Phe Lys Arg Met Gln Lys Ile Met Glu Glu Val Lys
112    115    120    125
114 Lys Ala Ser Gln Thr Val Arg Ile Gln Glu Glu Leu Ser Val Leu Leu
115    130    135    140
117 Asn Glu Asp Ile Val Leu Ser Ile Asp Ser Ser Ala Asp Tyr Ile Asp
118 145      150      155      160
120 Ala Val Ile Lys Val Leu Asp Val Leu Phe Lys Ile Ile Asn Val Arg
121      165      170      175
123 Ser
126 <210> SEQ ID NO: 3
127 <210> LENGTH: 846
128 <212> TYPE: DNA
129 <213> ORGANISM: Chlamydia trachomatis
131 <220> FEATURE:
132 <221> NAME/KEY: CDS
133 <222> LOCATION: (1) (846)
135 <400> SEQUENCE:
136 atg aat cgc atg att tct gat tgc gta tct cgc ata act ggc gat cga      16
137 Met Asn Arg Met Ile Cys Asp Cys Val Ser Arg Ile Thr Gly Asp Arg
138      1      5      10      15

```

RAW SEQUENCE LISTING

PATENT APPLICATION US/09/673,763

DATE 01/13/2001

TIME 10:34:38

Input Set A:\55928 seq.txt

Output Set N:\CRF3\01182001\1673763.raw

```

140 gtc aag aat att gtt ctg att gat gga ggc att gat cct cat tca tat 96
141 Val Lys Asn Ile Val Leu Ile Asp Gly Ala Ile Asp Pro His Ser Tyr
142      20      25      30
144 gag arg gtc aag ggt gat gaa ggg gaa atc ggc att agc cag ctg att 141
145 Glu Met Val Lys Gly Asp Glu Asp Arg Met Ala Met Ser Gln Leu Ile
146      35      40      45
148 ttt tgc aat gat tta ggt tta gag cat tca gct agt tta cgt aac cat 192
149 Phe Cys Asn Glu Leu Gly Leu Glu His Ser Ala Ser Leu Arg Lys His
150      50      55      60
152 cta gag ggt aac cca aca gtc gtt gat tta ggt aac cgt ttg ctt aac 240
153 Leu Glu Glu Lys Pro Lys Val Val Asp Leu Glu Glu Asp Leu Leu Asn
154      65      70      75
156 aac aac cgt ttt gat ctt ctg agt aca gaa gga ttc cct gac cca cat 288
157 Lys Asn Cys Phe Asp Leu Leu Ser Glu Glu Glu Phe Pro Asp Pro His
158      85      90      95
160 att tca acg gat atc aga gta tgg gat cct cat gta aac gag atg gct 336
161 Ile Trp Thr Asp Met Arg Val Trp Glu Ala Ala Val Lys Glu Met Ala
162      100      105      110
164 gcg gca tta att cca aca ttt cct caa taa gaa gag gal ttt cca gag 384
165 Ala Ala Leu Ile Gln Gln Phe Pro Gln Lys Glu Glu Asp Phe Gln Lys
166      115      120      125
168 aat gcg gat aag att tta tca gac att gcg aca cct gat cgt tgg gaa 432
169 Asn Ala Asp Glu Ile Leu Ser Glu Met Glu Glu Leu Asp Arg Trp Ala
170      130      135      140
172 gtc cgt tct ctg tct acg att cct gaa aca aat cgt tat tta gtc aca 480
173 Val Arg Ser Leu Ser Thr Ile Pro Glu Lys Asn Arg Tyr Leu Val Phe
174      145      150      155
176 ggc cac aat gcg ttc agt tac ttt act cgt aag tat cta tcc tct gat 528
177 Gly His Asn Ala Phe Ser Thr Phe Thr Arg Tyr Leu Ser Ser Asp
178      155      160      165
180 gcg cag aca gta tct gcg gaa tgg aca tgc cgt ttc att tct cca gaa 576
181 Ala Glu Arg Val Ser Gly Glu Trp Arg Ser Arg Cys Ile Ser Pro Glu
182      170      175      180
184 gga ttg tct cct gag gct cag att agt ctc cga gaa att atg cgt gaa 624
185 Gly Leu Ser Pro Glu Ala Gln Ile Ser Ile Arg Asp Ile Met Arg Val
186      185      190      195
188 gtc gag tat atc tct gca aac gat gta cca gtt gtc thr tta gag cat 672
189 Val Glu Trp Ile Ser Ala Asn Asp Val Glu Val Val Phe Leu Glu Asp
190      200      205      210
192 acg tta aat caa gat gct ttg aca aag att gtt tct tgc tct aag agc 720
193 Thr Leu Asn Gln Asp Ala Leu Arg Lys Ile Val Ser Cys Ser Lys Ser
194      215      220      225
196 gga caa aag att cgt ac tct aag tct cct tta tct agc gat aat gtc 768
197 Gly Gln Lys Ile Arg Leu Ala Lys Ser Pro Leu Tyr Ser Asp Asn Val
198      230      235      240
200 tgt gat aac tat ttt agc acg ttc cag cac aca gtt cgc aca att aca 816
201 Cys Asp Asn Tyr Phe Ser Thr Phe Gln His Asn Val Arg Phe Ile Thr
202      245      250      255
204 gaa gaa ttg gga ggg act gtt ctt gaa taa 840

```

RAW SEQUENCE LISTING
 PATENT APPLICATION US/09/673,763
 DATE: 01/13/2001
 TIME: 10:34:30

Input Seq: A:\55928 seq.txt
 Output Seq: N:\CRF3\01182001\I673763.raw

```

205 Glu Cln Ier Gly Gly Lys Val Leu Glu
206      275      280
209 <211> SEQ ID NO: 4
210 <211> LENGTH: 281
211 <212> TYPE: PRT
212 <213> ORGANISM: Chlamydia trachomatis
214 <400> SEQUENCE:
215 Met Asn Arg Met Ile Cys Asp Cys Val Ser Arg Ile Lys Gly Asp Ala
216      1      5      10      15
218 Val Lys Asn Ile Val Leu Ile Asp Gly Ala Ile Asp Pro His Ser Tyr
219      20      25      30
221 Glu Met Val Cys Gly Asp Cln Asp Ala Met Ala Met Ser Gly Leu Ile
222      35      40      45
224 Phe Cys Asn Gly Leu Gly Leu Glu His Ser Ala Ser Leu Arg Lys His
225      50      55      60
227 Leu Glu Gly Asn Pro Lys Val Val Asp Leu Gly Glu Arg Leu Leu Asn
228      65      70      75      80
230 Lys Asn Cys Phe Asp Leu Leu Ser Glu Glu Gly Phe Pro Asp Pro Asn
231      85      90      95
233 Ile Trp Thr Asp Met Arg Val Trp Gly Ala Ala Val Lys Glu Met Ala
234      100      105      110
236 Ala Ala Leu Ile Glu Gly Phe Pro Gly Tyr Glu Glu Asp Phe Gly Lys
237      115      120      125
239 Asn Ala Asp Glu Ile Leu Ser Cln Met Glu Glu Leu Asp Ala Trp Ala
240      130      135      140
242 Val Arg Ser Leu Ser Thr Ile Pro Glu Lys Asn Arg Tyr Leu Val Phe
243      145      150      155      160
245 Gly His Asn Ala Phe Ser Tyr Phe Thr Arg Arg Tyr Leu Ser Ser Asp
246      165      170      175
248 Ala Glu Arg Val Ser Gly Cln Trp Arg Ser Arg Cys Ile Ser Pro Glu
249      180      185      190
251 Gly Leu Ser Pro Gly Ala Glu Ile Ser Ile Arg Asp Ile Met Arg Val
252      195      200      205
254 Val Glu Tyr Ile Ser Ala Asp Asp Val Glu Val Val Phe Leu Glu Asp
255      210      215      220
257 Thr Leu Asn Gln Asp Ala Leu Arg Lys Ile Val Ser Cys Ser Lys Ser
258      225      230      235      240
260 Gly Glu Lys Ile Arg Leu Ala Lys Ser Pro Leu Tyr Ser Asp Asn Val
261      245      250      255
263 Cys Asp Asn Tyr Phe Ser Thr Phe Gly Asn Asn Val Arg Thr Ile Thr
264      260      265      270
266 Glu Glu Leu Gly Gly Thr Val Leu Glu
267      275      280
270 <216> SEQ ID NO: 5
271 <211> LENGTH: 861
272 <212> TYPE: PRT
273 <213> ORGANISM: Chlamydia trachomatis
275 <220> FEATURE
276 <221> NAME/KEY: CDS

```

RAW SEQUENCE LISTING

PATENT APPLICATION US/09/672,763

Date: 01/13/2001

Time: 10:30:38

Input Seq: A:\55928 seq.txt

Output Seq: N:\CRF3\01182001\I673763.raw

```

277 <222> LOCATION (1) (861)
279 <400> SEQUENCE
280 atg tct ggg ata acc att tca gga cgt ttc agc acc atg atc gca caa 48
281 Met Ser Val Ile Thr Ile Leu Ala Arg Ser Ser Thr Met Phe Ala Gln
282 1 5 10 15
284 tta cag cag aat tgg gag gga ctg ctg ttg aat aga gat aat gca att 96
285 Leu Glu Lys Asn Trp Glu Gly Leu Phe Leu Asn Arg Asp Asn Ala Ile
286 20 25 30
288 gct tgg ttc gta gag gtt ctt tgt gtt aat tat gat cac tca gag gtc 144
289 Ala Trp Ser Val Glu Asp Leu Cys Val Asn Tyr Asp His Ser Asp Val
290 35 40 45
292 tta tcc ccc att aat ttt ttc ctg cct gcc ggg gga atg gct cct ctt 192
293 Leu Cys His Ile Thr Phe Ser Leu Pro Ala Gly Ala Met Ala Ala Ile
294 50 55 60
296 att gcc cag aat aga gct ggt aac aat aat ttg ctt aag gct tct tta 240
297 Ile Gly Pro Asn Gly Ala Gly Lys Ser Thr Leu Leu Lys Ala Ser Leu
298 65 70 75 80
300 gga ctg aat cgt gct tct cct agc caa aac ttg ttc ttt ggt cag aga 288
301 Gly Leu Ile Arg Ala Ser Ser Cys Glu Ser Leu Phe Phe Gly Gln Arg
302 85 90 95
304 ttt tcc aag gca tat tat aga ata cct tat atg cct caa aga gga aat 336
305 Phe Ser Lys Ala His His Arg Ile Ala Tyr Met Pro Gln Arg Ala Ser
306 100 105 110
308 gtc gat tgg gat ttc tca atc act gtt ctt gat ctg gtg ttg atg ggg 384
309 Val Asp Trp Asp Phe Pro Met Thr Val Leu Asp Leu Val Leu Met Gly
310 115 120 125
312 tgt tcc ggc tat aca gga cta tgc aat cgt aat tcc act gat gat cgt 432
313 Cys Tyr Gly Tyr Lys Gly Leu Trp Asn Arg Ile Ser Thr Asp Asp Arg
314 130 135 140
316 cag aag gct atg cgt att tta gag ccc att agt ttg aac gct ttt gca 480
317 Glu Glu Ala Met Arg Ile Leu Glu Arg Val Gly Leu Glu Ala Phe Ala
318 145 150 155 160
320 aat cgt aca ata agt aag ctg tcc tcc aga gga caa caa cag aga gct ctt 528
321 Asn Arg Glu Ile Gly Lys Leu Ser Gly Tyr Glu Gln Gln Arg Ala Phe
322 165 170 175
324 tta cag cag tca tta arg caa aca gca gat tgg tat ctg atg gat gag 576
325 Leu Ala Arg Ser Leu Met Gln Lys Ala Asp Leu Tyr Leu Met Asp Glu
326 180 185 190
328 ctg ttc tct cgg atc gat atg gtc cct tat caa atg gtt gla gat gtt 624
329 Leu Phe Ser Ala Ile Asp Met Ala Ser Tyr Gln Met Val Val Asp Val
330 195 200 205
332 ttg caa ggg ctt aac agc gaa ggg aag cct att gtc gtc att cat cat 672
333 Leu Cys Glu Leu Lys Ser Gly Gly Lys Thr Ile Val Val Ile His His
334 210 215 220
336 gat ttg agt aat gtc cag aag ctt ctt gat cat cgg att tta tta aat 720
337 Asp Glu Ser Asn Val Arg Lys Leu Phe Asp His Val Ile Leu Leu Asn
338 225 230 235 240
340 aag cat ctt gtc tgc tct gga agc gla gaa gaa tgc ttg act aac gaa 768
341 Lys His Leu Val Cys Ser Gly Ser Val Glu Glu Cys Leu Thr Lys Glu

```

TARGET APPLICATION US/09/673,763

TIME 10 34 30

Output Set N:\CRF3\01132001\I673763.raw

file://C:\CRF3\Outhold\Vsr1673763.htm

COMBINED DECLARATION AND POWER OF ATTORNEY FOR PATENT APPLICATION

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,

I believe I am an original, first and joint inventor of the subject matter which is claimed and for which a patent is sought on the invention entitled CHLAMYDIA PROTEINS AND THEIR USES, the specification of which

- ☒ is attached hereto.
- ☐ was filed on _____ as Application No. _____.
- ☐ was described and claimed in PCT International Application No. _____, filed on _____.
- ☐ and was amended on _____ (if applicable).
- ☐ with amendments through _____ (if applicable).

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, Section 1.56. If this is a continuation-in-part application filed under the conditions specified in 35 U.S.C. § 120 which discloses and claims subject matter in addition to that disclosed in the prior copending application, I further acknowledge the duty to disclose material information as defined in 37 C.F.R. § 1.56 which occurred between the filing date of the prior application and the national or PCT international filing date of the continuation-in-part application.

I hereby claim foreign priority benefits under Title 35, United States Code, Section 119(a)-(d) of any foreign application(s) for patent or inventor's certificate or of any PCT International application(s) designating at least one country other than the United States of America listed below and have also identified below any foreign application(s) for patent or inventor's certificate or any PCT International application(s) designating at least one country other than the United States of America filed by me on the same subject matter having a filing date before that of the application(s) on which priority is claimed:

Prior Foreign Application(s)

Priority
Claimed

(Number)

(Country)

(Day/Month/Year Filed)

☐
Yes

☐
No

I hereby claim the benefit under Title 35, United States Code, Section 119(e) of any United States provisional application(s) listed below:

60/082,438

60/082,588

60/086,450

Application Number

20 April 1998

21 April 1998

22 May 1998

Filing Date

I hereby claim the benefit under Title 35, United States Code, Section 120 of any United States application(s) or Section 365(c) of any PCT International application(s) designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of Title 35, United States Code, Section 112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, Section 1.56(a) which occurred between the filing date of the prior application and the national or PCT International filing date of this application:

PCT/US99/08744

(Application No.)

20 April 1999

(Filing Date)

Published

(Status)

The undersigned hereby authorizes the U.S. attorney or agent named herein to accept and follow instructions from _____ as to any action to be taken in the Patent and Trademark Office regarding this application without direct communication between the U.S. attorney or agent and the undersigned. In the event of a change in the persons from whom instructions may be taken, the U.S. attorney or agent named herein will be so notified by the undersigned.

I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application, to file a corresponding international application, and to transact all business in the Patent and Trademark Office connected therewith:

☒ Customer Number

-OR-

☐ Registered practitioners listed below:
**24197**

Name	Reg. No.	Name	Reg. No.
BECKER, Mark L.	31,325	ORR, David E.	44,988
CALDWELL, Lisa M.	41,653	PETERSEN, David P.	28,106
DeGRANDIS, Paula A.	43,581	POLLEY, Richard J.	28,107
GEORGE, Samuel E.	44,119	RINEHART, Kyle B.	47,027
GIRARD, Michael P.	38,467	SCOTTI, Robert F.	39,830
HARDING, Tanya M.	42,630	SIEGEL, Susan Alpert	43,121
JAKUBEK, Joseph T.	34,190	SLATER, Stacey C.	36,011
JONES, Michael D.	41,879	STEPHENS Jr., Donald L.	34,022
KLARQUIST, Kenneth S.	16,445	STUART, John W.	24,540
KLITZKE II, Ramon A.	30,188	VANDENBERG, John D.	31,312
LEIGH, James S.	20,434	WHINSTON, Arthur L.	19,155
MAURER, Gregory L.	43,781	WIGHT, Stephen A.	37,759
NOONAN, William D.	30,878	WINN, Garth A.	33,220

Address all telephone calls to Tanya M. Harding, Ph.D. at telephone number (503) 226-7391.


Address all correspondence to:


KLARQUIST SPARKMAN CAMPBELL
LEIGH & WHINSTON, LLP
One World Trade Center, Suite 1600
121 S.W. Salmon Street
Portland, OR 97204-2988

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full Name of Sole or first Inventor:

Daniel D. Rockey


Inventor's Signature



10/12/00
Date


Residence: Corvallis, OR


Citizenship: United States of America

Post Office Address: 3625 NW Jackson Street
Corvallis, OR 97330

Full Name of Second Inventor:

John P. Bannantine


Inventor's Signature



October 5, 2000
Date

Residence: Ames, IA

Citizenship: United States of America

Post Office Address: 903 Yuma Ave.
Ames, IA 50014